

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 0 911 333 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
28.04.1999 Bulletin 1999/17

(51) Int Cl.⁶: **C07D 487/04, A61K 31/505**
// (C07D487/04, 239:00, 231:00)

(21) Application number: **98308177.9**

(22) Date of filing: **08.10.1998**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **Bell, Andrew Simon**
Sandwich, Kent CT13 9NJ (GB)
• **Terrett, Nicholas Kenneth, Dr.**
Sandwich, Kent CT13 9NJ (GB)


(30) Priority: **24.10.1997 GB 9722520**

(74) Representative: **Harding, Charles Thomas**
D. Young & Co.
21 New Fetter Lane
London EC4A 1DA (GB)

(71) Applicants:
• **PFIZER INC.**
New York, N.Y. 10017 (US)
Designated Contracting States:
BE CH DE DK ES FI FR GR IE IT LI LU NL PT AT CY
• **Pfizer Limited**
Sandwich Kent CT13 9NJ (GB)
Designated Contracting States:
GB

(54) **Pyrazolo[4,3-d]pyrimidine derivatives and pharmaceutical compositions containing them**

(57) Compounds are disclosed which are useful as PDE1 inhibitors. The compounds have the formula:



(I)

EP 0 911 333 A1

Description


[0001] The present invention relates to compounds, including *inter alia* pharmaceutical compositions comprising the same and methods for making the same.

[0002] In particular, the present invention relates to compounds that are capable of exhibiting inhibition of a phosphodiesterase (PDE) enzyme.

[0003] More in particular, the present invention relates to compounds that are capable of exhibiting at least inhibition of a phosphodiesterase type 1 (PDE1) enzyme - i.e. the compounds are capable of acting as inhibitors of PDE1. Some of these compounds are also capable of exhibiting inhibition of other types of PDE enzymes - such as a phosphodiesterase type 5 (PDE5) enzyme.

[0004] By way of background information, EP-A-0201188 discloses certain 5-substituted pyrazolo[4,3-d]pyrimidin-7-ones and suggests their use for the treatment of cardiovascular disorders, such as heart failure or cardiac insufficiency. EP-A-0201188 also suggests the use of those 5-substituted pyrazolo[4,3-d]pyrimidin-7-ones to inhibit PDE.

[0005] In particular, Example 1 of EP-A-0201188 discloses the following 5-substituted pyrazolo[4,3-d]pyrimidin-7-one:



[0006] For ease of reference this compound will be referred to as the '188 Compound. Like the '188 compound, all of the compounds of EP-A-0201188 have a methyl group attached at the 3 position of the pyrazolo[4,3-d] pyrimidine ring system.


[0007] We have found that the '188 compound is at least a weak PDE1 inhibitor.

[0008] Furthermore, as there is now a body of evidence associating PDE1 with a number of diseases, e.g. stroke, dementia, memory enhancement, atherosclerosis, urge incontinence, hypertension, angina pectoris, congestive heart failure, myocardial infarction and restenosis, so there is a need to have more potent PDE1 inhibitors.

[0009] There is also a need to have more selective PDE inhibitors, in particular PDE1 inhibitors.

[0010] The present invention seeks to provide compounds that are useful as PDE1 inhibitors, including pharmaceutical compositions comprising the same and methods for making the same.

[0011] According to a first aspect of the present invention there is provided a compound of the formula (I)



wherein

R_a is C₂-C₆ alkyl;


R₁ is H or C₁-C₄ alkyl;

each of R₂ and R₃ is independently selected from H and C₁-C₄ alkyl, or R₂ is H or C₁-C₄ alkyl and R₃ is OH, C₂-C₄ alkanoyloxy or fluoro, or

R₂ and R₃ when taken together represent C₂-C₆ alkylene, or

R₂ and R₃ when taken together with the carbon atom to which they are attached represent a carbonyl group;

Ar is either (a)



wherein each of R₄, R₅ and R₆ is independently selected from

H,
 C₁-C₄ alkyl,
 C₁-C₄ alkoxy,
 C₁-C₄ alkoxy-Z-,
 halo,
 halo(C₁-C₄)alkyl,
 phenoxy, optionally substituted by up to three substituents each of which substituent is independently selected from halo, C₁-C₄ alkyl, and C₁-C₄ alkoxy,
 nitro,
 hydroxy,
 hydroxy-Z-,
 C₂-C₄ alkanoyl,
 amino,
 amino-Z-,
 (C₁-C₄ alkyl)NH,
 (C₁-C₄ alkyl)₂N-,
 (C₁-C₄ alkyl)NH-Z-,
 (C₁-C₄ alkyl)₂N-Z-,
 -COOH,
 -Z-COOH,
 -COO(C₁-C₄ alkyl),
 -Z-COO(C₁-C₄ alkyl),
 C₁-C₄ alkanesulphonamido,
 C₁-C₄ alkanesulphonamido-Z-,
 halo(C₁-C₄)alkanesulphonamido,
 halo(C₁-C₄)alkanesulphonamido-Z-,
 C₁-C₄ alkanamido,
 C₁-C₄ alkanamido-Z-,
 HOOC-Z-NH-,
 HOOC-Z-NH-Z-,
 (C₁-C₄ alkyl)OOC-Z-NH-,
 (C₁-C₄ alkyl)OOC-Z-NH-Z-,
 C₁-C₄ alkyl-NH-SO₂-NH-,
 C₁-C₄ alkyl-NH-SO₂-NH-Z-,

(C₁-C₄ alkyl)₂-N-SO₂-NH-,
 (C₁-C₄ alkyl)₂-N-SO₂-NH-Z-,
 C₁-C₄ alkoxy CH=CH-Z-CONH-,
 C₁-C₄ alkoxy CH=CHCONH-

C₁-C₄ alkyl-SO₂-N(C₁-C₄ alkyl)-,
 C₁-C₄ alkyl-SO₂-N(C₁-C₄ alkyl)-Z-,

(C₁-C₄ alkyl)NH-Z-SO₂-NH-,

(C₁-C₄ alkyl)₂N-Z-SO₂-NH-,

(C₁-C₄ alkyl)NH-Z-SO₂-NH-Z-,

(C₁-C₄ alkyl)₂N-Z-SO₂-NH-Z-,

benzenesulphonamido, optionally ring substituted by up to three substituents each of which is independently selected from halo, C₁-C₄ alkyl, and C₁-C₄ alkoxy,

C₁-C₄ alkanoyl-N(C₁-C₄ alkyl)-,

C₁-C₄ alkanoyl-N(C₁-C₄ alkyl)-Z-,

C₁-C₄ alkoxycarbonyl-CH(CH₂OH)NHSO₂-,

-SO₃H,

-SO₂NH₂,

H₂NOC-CH(CH₂OH)-NHSO₂-,

HOOC-Z-O-, and

(C₁-C₄ alkyl)OOC-Z-O-,

or optionally one of R₄, R₅ and R₆ is a G-Het group and wherein the others of R₄, R₅ and R₆ are independently selected from the R₄, R₅ and R₆ substituents listed above;

Z is C₁-C₄ alkylene,

G is a direct link, Z, O, -SO₂NH-, SO₂, or -Z-N(C₁-C₄ alkyl)SO₂-,

Het is a 5- or 6-membered heterocyclic group containing 1, 2, 3 or 4 nitrogen heteroatoms; or 1 or 2 nitrogen heteroatoms and 1 sulphur heteroatom or 1 oxygen heteroatom; or the heterocyclic group is furanyl or thiophenyl; wherein the Het group is saturated or partially or fully unsaturated and optionally substituted by up to 3 substituents, wherein each substituent is independently selected from C₁-C₄ alkyl, oxo, hydroxy, halo, and halo(C₁-C₄) alkyl;

or (b) any one of the following bicyclic groups:

benzodioxolanyl,

benzodioxanyl,

benzimidazolyl,

quinoliny,

indolyl,

quinazolinyl,

isoquinoliny,

benzotriazolyl,

benzofuranyl,

benzothiophenyl,

quinoxaliny,

or

phthaliziny,

wherein said bicyclic Ar groups are linked to the neighbouring -C(R₂R₃)- group *via* the benzo ring portion,

and wherein the heterocyclic portion of said bicyclic Ar group is optionally partially or fully saturated, said group being optionally substituted by one or more of C₁-C₄ alkyl, halo, hydroxy, oxo, amino, and C₁-C₄ alkoxy;


or a pharmaceutically acceptable salt of the compound, or a pharmaceutically acceptable solvate of the compound or the salt.

[0012] This and some of the other aspects of the present invention, as well as some preferred embodiments of the present invention, are presented in the accompanying claims.

[0013] It will also be appreciated that what is to be claimed includes the following:


- (i) a compound of the formula (I) or a pharmaceutically acceptable salt thereof;
- (ii) one or more processes for the preparation of a compound of the formula (I) or a pharmaceutically acceptable salt thereof;
- (iii) novel intermediates for use in any one of those processes;
- (iv) a pharmaceutical composition comprising a compound of the formula (I), or a pharmaceutically acceptable salt thereof, admixed with a pharmaceutically acceptable diluent, carrier or excipient;
- (v) a compound of the formula (I), or a pharmaceutically acceptable salt or composition thereof, for use as a medicament;
- (vi) the use of a compound of the formula (I), or of a pharmaceutically acceptable salt or composition thereof, for the manufacture of a medicament for the treatment of conditions capable of being treated by the inhibition of PDE enzymes;
- (vii) use as in (vi) wherein the medicament is for use as an inhibitor for PDE1;
- (viii) a method of treatment of a subject (e.g. a mammal) in need of same, which method comprises administering to the subject an effective amount of a compound of the formula (I) or a pharmaceutically acceptable salt or composition thereof, and wherein the compound, salt or composition produces an inhibitory effect against a PDE; and
- (ix) a method as in (viii) wherein the compound, salt or composition produces an inhibitory effect against PDE1.

[0014] By way of example, a preferred process according to one embodiment of the present invention for preparing compounds according to the present invention is presented by the following scheme:



wherein each of the groups are as defined above.


[0015] By way of further example, a preferred process according to another embodiment of the present invention for preparing compounds according to the present invention is presented by the following scheme:



wherein each of the groups are as defined above, and wherein R_7 is H or C_{1-4} alkyl, and wherein X is C_{1-4} alkyl, halo (C_{1-4})alkyl, or optionally substituted phenyl.


[0016] By way of further example, a preferred process according to another embodiment of the present invention for

preparing compounds according to the present invention is presented by the following scheme:




wherein each of the groups are as defined above.

[0017] By way of example, a preferred intermediate for one embodiment of the present invention is




wherein each of the groups are as defined above.

[0018] By way of further example, a preferred intermediate for another embodiment of the present invention is



wherein each of the groups are as defined above.

[0019] By way of further example, a preferred intermediate for another embodiment of the present invention is



wherein each of the groups are as defined above.

[0020] A key advantage of the present invention is that it provides compounds, and compositions comprising the same, that are useful as PDE1 inhibitors.

[0021] Another key advantage of the compounds of the present invention is that some are selective PDE inhibitors, in particular selective PDE1 inhibitors.

[0022] As indicated above, the compounds of the present invention are of the general formula (I). We have surprisingly

found that these compounds are effective as PDE1 inhibitors and at low concentrations. This result is surprising because the compounds of EP-A-0201188, such as the '188 compound, are not as effective at such low concentrations. This highly surprising result is borne out by the experimental data presented in the experimental section (*infra*).

[0023] The compounds of the present invention may exist in hydrated or solvated forms.

[0024] Alkyl and alkylene groups, when present in any one of the above-defined groups for the compounds of the formula (I), may be linear or branched.

[0025] The term "halo" as used herein means means F, Cl, Br or I.

[0026] The pharmaceutically acceptable salts of the compounds of the formula (I) include suitable acid addition or base salts thereof. For a review on suitable pharmaceutical salts see Berge *et al*, J Pharm Sci, **66**, 1-19 (1977).

[0027] By way of example, suitable acid addition salts are formed from acids which form non-toxic salts. Suitable examples of such salts are the hydrochloride, hydrobromide, hydroiodide, sulphate, bisulphate, phosphate, hydrogen phosphate, acetate, maleate, fumarate, lactate, tartrate, citrate, gluconate, benzoate, methanesulphonate, benzenesulphonate and *p*-toluenesulphonate salts.

[0028] Also by way of example, suitable base salts are formed from bases which form non-toxic salts. Suitable examples thereof are the aluminium, calcium, lithium, magnesium, potassium, sodium, zinc, N-benzyl-N-(2-phenylethyl) amine, 1-adamantylamine and diethanolamine salts.

[0029] Compounds of the present invention may contain one or more asymmetric carbon atoms and/or one or more non-aromatic carbon-carbon double bonds and may therefore exist in two or more stereoisomeric forms. Thus, the present invention also provides both the individual stereoisomers of the compounds of the formula (I), as well as mixtures thereof, including compositions comprising the same. Separation of diastereoisomers or *cis* and *trans* isomers may be achieved by conventional techniques, e.g. by fractional crystallisation, chromatography or HPLC of a stereoisomeric mixture of a compound of the formula (I) or a suitable salt or derivative thereof. An individual enantiomer of a compound of the formula (I) may also be prepared from a corresponding optically pure intermediate or by resolution, such as by HPLC of a racemate using a suitable chiral support or by fractional crystallisation of the diastereoisomeric salts formed by reaction of a racemate with a suitable optically active acid or base.

[0030] As mentioned above, the present invention also covers pharmaceutical compositions comprising the compounds of the general formula (I). In this regard, and in particular for human therapy, even though the compounds of the present invention (including their pharmaceutically acceptable salts and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutical carrier, excipient or diluent selected with regard to the intended route of administration and standard pharmaceutical practice.

[0031] By way of example, in the pharmaceutical compositions of the present invention, the compounds of the present invention may be admixed with any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), or solubilising agent(s).

[0032] In general, a therapeutically effective daily oral or intravenous dose of the compounds of formula (I) and their salts is likely to range from 0.01 to 50 mg/kg body weight of the subject to be treated, preferably 0.1 to 20 mg/kg. The compounds of the formula (I) and their salts may also be administered by intravenous infusion, at a dose which is likely to range from 0.001-10 mg/kg/hr.

[0033] Tablets or capsules of the compounds may be administered singly or two or more at a time, as appropriate. It is also possible to administer the compounds in sustained release formulations.

[0034] Typically, the physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. The above dosages are exemplary of the average case. There can, of course, be individual instances where higher or lower dosage ranges are merited, and such are within the scope of this invention.

[0035] Alternatively, the compounds of the general formula (I) can be administered by inhalation or in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. An alternative means of transdermal administration is by use of a skin patch. For example, they can be incorporated into a cream consisting of an aqueous emulsion of polyethylene glycols or liquid paraffin. They can also be incorporated, at a concentration of between 1 and 10% by weight, into an ointment consisting of a white wax or white soft paraffin base together with such stabilisers and preservatives as may be required.

[0036] For some applications, preferably the compositions are administered orally in the form of tablets containing excipients such as starch or lactose, or in capsules or ovules either alone or in admixture with excipients, or in the form of elixirs, solutions or suspensions containing flavouring or colouring agents.

[0037] The compositions (as well as the compounds alone) can also be injected parenterally, for example intracavernosally, intravenously, intramuscularly or subcutaneously. In this case, the compositions will comprise a suitable carrier or diluent.

[0038] For parenteral administration, the compositions are best used in the form of a sterile aqueous solution which may contain other substances, for example enough salts or monosaccharides to make the solution isotonic with blood.

[0039] For buccal or sublingual administration the compositions may be administered in the form of tablets or loz-

enges which can be formulated in a conventional manner.

[0040] For oral, parenteral, buccal and sublingual administration to subjects (such as patients), the daily dosage level of the compounds of the present invention and their pharmaceutically acceptable salts and solvates may typically be from 10 to 500 mg (in single or divided doses). Thus, and by way of example, tablets or capsules may contain from 5 to 100 mg of active compound for administration singly, or two or more at a time, as appropriate. As indicated above, the physician will determine the actual dosage which will be most suitable for an individual patient and it will vary with the age, weight and response of the particular patient. It is to be noted that whilst the above-mentioned dosages are exemplary of the average case there can, of course, be individual instances where higher or lower dosage ranges are merited and such dose ranges are within the scope of this invention.

[0041] Generally, in humans, oral administration of the compounds of the invention is the preferred route, being the most convenient and, for example in male erectile dysfunction (MED), avoiding the well-known disadvantages associated with intracavernosal (i.c.) administration. A preferred oral dosing regimen in MED for a typical man is from 25 to 100 mg of compound when required. In circumstances where the recipient suffers from a swallowing disorder or from impairment of drug absorption after oral administration, the drug may be administered parenterally, e.g. sublingually or buccally.

[0042] For veterinary use, a compound of the present invention or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate of either entity, is typically administered as a suitably acceptable formulation in accordance with normal veterinary practice and the veterinary surgeon will determine the dosing regimen and route of administration which will be most appropriate for a particular animal. However, as with human treatment, it may be possible to administer the compound alone for veterinary treatments.

[0043] Thus the invention provides a pharmaceutical composition comprising a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of either entity, together with a pharmaceutically acceptable diluent, excipient or carrier.

[0044] The present invention also provides a veterinary formulation comprising a compound of the present invention, or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate of either entity, together with a veterinarily acceptable diluent, excipient or carrier.

[0045] The invention further provides a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of either entity, or a pharmaceutical composition containing any of the foregoing, for use as a human medicament.

[0046] In addition, the present invention provides a compound of the present invention, or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate of either entity, or a veterinary formulation containing any of the foregoing, for use as an animal medicament.

[0047] In yet another aspect, the invention provides the use of a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of either entity, in the manufacture of a medicament for administration to a human for the treatment of a medical condition capable of being treated by the inhibition of PDE1 activity.

[0048] The present invention also provides the use of a compound of the present invention, or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate of either entity, in the manufacture of an animal medicament for the treatment of a medical condition capable of being treated by the inhibition of PDE1 activity.

[0049] In yet another aspect, the invention provides the use of a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate of either entity, in the manufacture of a medicament for administration to either a human or an animal wherein the medicament is for use as an inhibitor of PDE1.

[0050] Moreover, the present invention provides the use of a compound of the present invention, or a pharmaceutically acceptable salt thereof, or a pharmaceutically acceptable solvate containing either entity, in the manufacture of a human medicament for the treatment of any one or more of male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.

[0051] The present invention also provides the use of a compound of the present invention, or a veterinarily acceptable salt thereof, or a veterinarily acceptable solvate containing either entity, in the manufacture of an animal medicament for the treatment of any one or more of male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.

[0052] Additionally, the present invention provides a method of treating a medical condition for which a PDE1 inhibitor is required, in a mammal (including a human being), which comprises administering to said mammal a therapeutically

effective amount of a compound of the present invention, or a pharmaceutically or veterinarily acceptable salt thereof, or a pharmaceutically or veterinarily acceptable solvate of either entity, or a pharmaceutical composition or veterinary formulation containing any of the foregoing.

[0053] Still further, the present invention provides a method of treating any one or more of male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility in a mammal (including a human being), which comprises administering to said mammal a therapeutically effective amount of a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt thereof, or a pharmaceutically or veterinarily acceptable solvate of either entity, or a pharmaceutical composition or veterinary formulation containing any of the foregoing.

[0054] References to treatment (as well as treating) include any one or more of curative, palliative or prophylactic treatment of a disease or condition.

[0055] The compounds of the formula (I) can be prepared by novel routes or, alternatively, by conventional routes.

[0056] The compounds of the present invention may be prepared by any one of the synthesis processes presented in the Route Section (*infra*), or by any one of the more specific synthesis protocols presented in the Examples Section (*infra*) - which are presented as either Preparations or Examples. The present invention also encompasses any one or more of these processes, including any of the steps thereof, in addition to any novel intermediate(s) obtained therefrom or used therein.


[0057] The general syntheses of the compounds of the present invention are now presented in the following Route Section.

[0058] It is to be noted that in the following Route Section that a propyl group has been used as an example of a suitable R_a group. Naturally, compounds with other R_a groups may be used to prepare compounds of the present invention. Likewise, suitable substituents other than those presented for R_1 etc. may be used.

ROUTE SECTION

Route A

[0059] A compound of formula (I) may be obtained from a compound of formula (II) wherein




R_{1-3} and Ar are as previously defined for formula (I), by the application of known cyclisation methods for pyrimidinone ring formation. Thus, for example, the cyclisation may be effected by the treatment of (II) with a base such as sodium or potassium hydroxide, or sodium or potassium carbonate, optionally in the presence of hydrogen peroxide, in a C_1 to C_4 alcohol-water medium, at from about 50°C to the reflux temperature of the reaction mixture.

[0060] The cyclisation may also be mediated by a sodium or potassium C_1 to C_5 alkoxide, in a C_1 to C_4 alcohol solvent, at from about 50°C to the reflux temperature of the reaction mixture.

[0061] Alternative cyclisation procedures involve the treatment of (II) with either polyphosphoric acid at from about 130°C to about 150°C or with anhydrous zinc chloride at from about 200°C to about 220°C .

[0062] In certain examples, the Ar group contains substituents which are chemically reactive under the cyclisation conditions and further reaction takes place under the standard conditions e.g. an alkyl bromide may hydrolyse to an alcohol and a trifluoromethyl group or an ester group may be converted to a carboxylic acid.


[0063] By way of example, a preferred embodiment of the above-mentioned route is as follows:



15 wherein the each of the groups are as defined above.

Route B

20 [0064] Compounds of the formula (I) in which R_4 is NH_2 can be prepared from the corresponding nitrobenzene by a reductive method.



wherein each of the other groups are as defined above.

35 [0065] In a typical procedure the reduction is carried out by catalytic hydrogenation e.g. using either a heterogeneous catalyst such as palladium, palladium- or rhodium-on-carbon, Raney nickel, or a homogeneous catalyst e.g. tris(triphenylphosphine)chlororhodium, in a suitable organic solvent e.g. industrial methylated spirit or ethyl acetate. The reaction is preferably carried out at from room temperature to the reflux temperature of the solvent and a pressure of from 1 to 5 atmospheres (100-500kPa).


40 [0066] The reaction can also be carried out using an excess of an electron transfer reducing agent such as tin (II) chloride in a suitable solvent such as a C_1 to C_4 alcohol e.g. ethanol, at the reflux temperature of the reaction mixture.

Route C

45 [0067] Compounds of the formula (I) in which R_4 is a group of the formula $-NR_7SO_2X$, wherein R_7 and X are as defined above, can be prepared by reaction of a compound of formula (I), where R_4 is NHR_7 with an appropriate alkyl sulphonyl halide.

50

55



Here, each of the groups are as defined above.


15 **[0068]** Similarly, compounds of the formula (I) in which R_4 is NR_7COX can be prepared by acylation of the same starting material with an appropriate acid chloride or anhydride. Compounds in which R_4 is NR_7CHO may also be prepared from a mixed anhydride such as $(C_1-C_4 \text{ alkylCO})OCHO$.

20 **[0069]** The reaction may be carried out in a suitable inert solvent such as dichloromethane in the presence of an acid acceptor such as triethylamine or pyridine (which can also be used as the solvent), at a temperature of from $0^\circ C$ to the reflux temperature of the solvent, preferably at room temperature.

[0070] Compounds of the formula (I) in which R_4 is a group of the formula $-NR_7CHO$ can also be prepared using a formyl transfer agent such as formyl- or 1,2-diformylhydrazine. In this case the reaction is preferentially carried out in the absence of solvent at the reflux temperature of the formylating agent.

25 Route D

[0071] Compounds of the formula (I) in which R_4 is a group of the formula $(C_{1-4} \text{ alkyl})R_7N-$ may be prepared by reduction of compounds of the formula (I) in which R_4 is $NR_7CO(C_{1-3} \text{ alkyl})$, wherein R_7 is H or $(C_{1-4} \text{ alkyl})$.




Here, each of the groups are as defined above.

[0072] The reducing agent is selected from reagents such as lithium aluminium hydride and diborane and is preferentially carried out in an inert solvent such as tetrahydrofuran at a temperature from room temperature to the reflux temperature of the solvent.

45 **[0073]** Alternatively, the products may be prepared by reduction of an imine (R_4 is $N=CH-(C_{1-3} \text{ alkyl})$) or iminium ion (R_4 is $[NR_7=CH-(C_{1-3} \text{ alkyl})]^+$) which may optionally be isolated. The reducing agent may be sodium acetoxyborohydride or sodium cyanohydride. The reaction may also be carried out using catalytic hydrogenation using a catalyst such as palladium-on-charcoal.


50 Route E

[0074] Compounds of the formula (I) in which R_4 is nitro can be prepared by nitration of the corresponding benzene derivative. The reaction is preferentially carried out using mixtures of concentrated nitric and sulphuric acids at a temperature from $0^\circ C$ to $100^\circ C$.



Here, each of the groups are as defined above

15 **[0075]** Similarly, compounds in which R_4 is chlorosulphonyl can be prepared by chlorosulphonylation of the corresponding benzene derivative. The reaction is preferentially carried out using chlorosulphonic acid as the solvent at a temperature from 0°C to 100°C.



30 With this route you may get a mixture of regio-isomers.

Route F

35 **[0076]** When Ar is a bicyclic group, a hetero-ring fused to the benzo portion can be formed using conventional ring forming reactions. For example, when the fused ring is a pyridone, the ring is formed by treatment of the corresponding b-ethoxypropenamamide with a strong acid such as sulphuric or hydrochloric acid.




Here, each of the groups are as defined above

Route G

50 **[0077]** Compounds of the formula (I) in which R_4 is a sulphamido or aminoalkanesulphonamido group can be prepared by treatment of the corresponding sulphamoyl halide or haloalkanesulphonamide with an excess of the amine in an aqueous or alcoholic solvent at a temperature of from room temperature to the reflux temperature of the solvent.

55 By way of example:




Here, each of the groups are as defined above, and m and n are independently selected from 0 and 1.

15 [0078] A compound of formula (A) where n is 1 and m is 0 or 1 can be cyclised to give the corresponding cyclic sultam. The reaction is carried out using a strong base such as sodium hydride in an inert solvent such as dimethylformamide at a temperature from 0°C to room temperature.

Route H


20 [0079] Compounds of the formula (I) in which R₄ is an aminosulphonyl group can be prepared from the corresponding chlorosulphonyl derivative -SO₂Cl by treatment with an appropriate amine, optionally in excess, in an aqueous or alcoholic solvent, at a temperature of from room temperature to the reflux temperature of the solvent. For example:



[0080] Similarly, when the chlorosulphonyl derivative is treated with aqueous alkali such as sodium hydroxide, a compound of the formula (I) in which R₄ is a group of the formula-SO₃H can be obtained.

Route I

40 [0081] Compounds of the formula (I) in which R₄ is a -G-Het group wherein G is a direct link and Het is attached to the adjacent phenyl ring by a nitrogen atom e.g. imidazol-1-yl can be prepared from the corresponding halophenyl derivative, where halo is preferably bromo or iodo, and the heterocycle. The reaction is preferably carried out in the presence of a base such as potassium carbonate, and a copper catalyst, preferably copper bronze. The reaction can be carried out in a high boiling solvent such as dimethylformamide, dimethylacetamide or N-methyl-2-pyrrolidinone at the reflux temperature of the solvent, or alternatively may be carried out without solvent, at the melt temperature of the mixture.




Route J


[0082] Compounds of the formula (I) in which the groups R_3 to R_6 contain a hydroxy or carboxyl function can be prepared from the corresponding ether or ester under conventional hydrolytic conditions. Ethers are preferably hydrolysed under strongly acidic conditions such as using concentrated hydrobromic acid at a temperature of between 100°C and 150°C. Ester hydrolysis is preferably carried out under basic conditions, for example using sodium or potassium hydroxide as base, optionally in the presence of hydrogen peroxide, in water or an alcoholic solvent such as ethanol. The reaction is carried out at from room temperature to the reflux temperature of the solvent.

Route K

[0083] Compounds of the formula (I) in which the groups R_2 and R_3 together form a carbonyl function are preferably prepared from the corresponding secondary alcohol by an oxidative method. The preferred oxidant is pyridinium chlorochromate and the reaction is preferably carried out in an inert solvent such as dichloromethane at room temperature.




[0084] An intermediate of the formula (II) is prepared by reaction of a carboxylic acid chloride, which may be derived by treatment of the corresponding carboxylic acid with oxalyl chloride in dichloromethane in the presence of a catalytic quantity of dimethylformamide, with an aminopyrazole derivative of the formula (III). The preparation of (III) is conventional - for example see the teachings of US-A-5,272,147.



[0085] The reaction is carried out in a suitable inert solvent such as dichloromethane in the presence of an acid acceptor such as triethylamine or in a basic solvent such as pyridine, at a temperature of from 0°C to the reflux temperature of the solvent, preferably at room temperature.

[0086] Interconversion of functional groups may also be carried out using a compound of the formula (II). Thus, for example, when R_4 is bromomethyl, the halide can be displaced by an appropriate amine preferably used in excess. Similarly the bromomethyl derivative can be reacted with an alcohol to provide an ether derivative, preferably using a metal salt such as silver nitrate. The alcohol is generally used as solvent and the reaction is preferably carried out at room temperature. For example:



15 [0087] The present invention will now be discussed only by way of further examples. The following Examples Section provides illustrations of the preparation of the compounds (I). The following Preparations Section provides illustrations of the preparation of *inter alia* novel starting materials.

[0088] In these sections, the ¹H nuclear magnetic resonance (NMR) spectra were recorded using either a Varian Unity 300 or a Varian Inova 400 spectrometer and were in all cases consistent with the proposed structures. Characteristic chemical shifts (δ) are given in parts-per-million downfield from tetramethylsilane using conventional abbreviations for designation of major peaks: e.g. s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; b, broad.

[0089] The mass spectra (m/z) were recorded using a Fisons Instruments Trio mass spectrometer in the thermospray ionisation mode. In the following sections, room temperature means 20 to 25°C.


[0090] In the following examples, propyl means n-propyl unless otherwise stated.

25 SYNTHESIS EXAMPLES

Example 1

30 5-(4-bromobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0091]



50 [0092] A 30% w/w solution of hydrogen peroxide (27 ml, 0.238mol) was added to a solution of sodium hydroxide (7.24g, 0.181mol) in water (400ml). A solution of N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-bromophenyl)acetamide (29.0g, 0.076mol) in ethanol (350ml) was then added, and the reaction stirred at reflux for 3 hours.


[0093] On cooling, the solution was diluted with water (200ml) and acidified to pH5 with 2N aqueous hydrochloric acid solution. The resulting white precipitate was extracted into dichloromethane (2x250ml). The organic extracts were dried (MgSO₄), filtered and concentrated under reduced pressure to give the title compound as a colourless solid. Crystallisation from acetonitrile gave colourless needles (19.33g), m.p. 193-194 °C.

55 [0094] Found: C, 53.09; H, 4.72; N, 15.41; C₁₆H₁₇BrN₄O requires C, 53.19; H, 4.74; N, 15.51%.

[0095] ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.85 (s, 2H), 4.10 (s, 3H), 7.25 (d, 2H), 7.45 (d, 2H), 12.30 (s, 1H) ppm.



Examples 2 to 24



[0096] The compounds of the following tabulated Examples of the general formula:-







were prepared from the corresponding carboxamide using similar methods to that used in Example 1.



Ex. No.	R ₁	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
2	-CH ₃		<p>Found: C, 68.24; H, 6.47; N, 20.04%. C₁₆H₁₈N₄O, requires C, 68.06; H, 6.43; N, 19.85%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.90 (s, 2H), 4.10 (s, 3H), 7.20-7.40 (m, 5H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 200-202°C.</p> <p>Crystallisation solvent: ethanol.</p>
3	-CH ₂ CH ₃		<p>Found: C, 68.58; H, 6.80; N, 18.82%. C₁₇H₂₀N₄O, requires C, 68.89; H, 6.80; N, 18.91%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.55 (t, 3H), 1.90 (m, 2H), 2.95 (t, 2H), 4.10 (s, 2H), 4.65 (q, 2H), 7.30-7.50 (m, 5H), 10.00 (s, 1H) ppm</p> <p>Melting point: 171-173°C.</p> <p>Crystallisation solvent: methanol.</p>



4	-CH ₃		<p>Found: C, 68.85; H, 6.79; N, 18.87%. C₁₇H₂₀N₄O, requires C, 68.89; H, 6.80; N, 18.91%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.75 (d, 3H), 1.90 (m, 2H), 2.95 (t, 2H), 4.15 (q, 1H), 4.25 (s, 3H), 7.30-7.40 (m, 5H), 9.15 (s, 1H) ppm</p> <p>Melting point: 154-157°C.</p> <p>Crystallisation solvent: hexane.</p>
5	-CH ₃		<p>Found: C, 65.00; H, 6.26; N, 17.58%. C₁₇H₂₀N₄O₂, requires C, 65.36; H, 6.45; N, 17.94%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.95 (t, 2H), 3.83 (s, 3H), 4.03 (s, 2H), 4.23 (s, 3H), 6.90 (d, 2H), 7.30 (d, 2H), 9.55 (s, 1H) ppm.</p> <p>Melting point: 195-198°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>



6	-CH ₃		<p>Found: C, 62.30; H, 5.46; N, 17.02%. C₁₇H₁₈N₄O₃, requires C, 62.56; H, 5.56; N, 17.17%. ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.75 (s, 2H), 4.10 (s, 3H), 5.95 (s, 2H), 6.70-6.95 (m, 3H), 12.15 (s, 1H) ppm. Melting point: 204-206°C. Crystallisation solvent: ethanol.</p>
7	-CH ₃		<p>Found: C, 70.38; H, 5.85; N, 14.96%. C₂₂H₂₂N₄O₂, requires C, 70.57; H, 5.92; N, 14.96%. ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.95 (t, 2H), 4.05 (s, 2H), 4.25 (s, 3H), 7.00 (m, 4H), 7.15 (m, 1H), 7.35 (m, 4H), 9.75 (s, 1H) ppm. Melting point: 167-168°C. Crystallisation solvent: acetonitrile.</p>



8	-CH ₃		<p>Found: C, 66.58; H, 7.26; N, 17.73%. C₂₂H₂₉N₃O₂, requires C, 66.81; H, 7.39; N, 17.71%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 4H), 2.00 (br.m, 2H), 2.35 (br.m, 5H), 2.70 (br.m, 2H), 2.90 (t, 2H), 4.00 (s, 2H), 4.25 (s, 3H) 4.35(br.m, 1H), 6.90 (d, 2H), 7.25 (d, 2H), 9.30 (s, 1H) ppm.</p> <p>Melting point: 194-195°C.</p> <p>Crystallisation solvent: acetonitrile.</p>
9	-CH ₃		<p>Found: C, 58.54; H, 5.28; N, 21.93 %. C₁₆H₁₇N₃O₃, requires C, 58.70; H, 5.23; N, 21.40%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.75 (t, 3H), 1.55 (m, 2H), 2.55 (t, 2H), 4.05 (s, 3H), 4.35 (s, 2H), 7.55 (m, 2H), 7.70 (m, 1H), 8.05 (d, 1H) 12.35 (s, 1H) ppm.</p> <p>Melting point: 217-222°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>



10	-CH ₃		<p>Found: C, 58.44; H, 5.24; N, 21.30%. C₁₆H₁₇N₃O₃, requires C, 58.70; H, 5.23; N, 21.40%. ¹H-NMR (DMSO-d₆): δ = 0.85 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 4.05 (s, 5H), 7.60 (t, 1H), 7.80 (d, 1H), 8.10 (d, 1H), 8.25 (s, 1H), 12.35 (s, 1H) ppm. Melting point: 230-233°C. Crystallisation solvent: ethyl acetate/methanol.</p>
11	-CH ₃		<p>Found: C, 58.46; H, 5.03; N, 21.08%. C₁₆H₁₇N₃O₃, requires C, 58.70; H, 5.23; N, 21.40%. ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.95 (t, 2H), 4.20 (s, 2H), 4.30 (s, 3H), 7.60 (d, 2H), 8.20 (d, 2H), 11.35 (s, 1H) ppm. Melting point: 225-226 °C. Crystallisation solvent: ethyl acetate.</p>




12	-CH ₃		<p>Found: C, 53.07; H, 4.76; N, 15.15%. C₁₆H₁₇BrN₄O, requires C, 53.19; H, 4.74; N, 15.51%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.60 (m, 2H), 2.60 (t, 2H), 4.05 (s, 2H), 4.10 (s, 3H), 7.20 (m, 2H), 7.35 (d, 1H), 7.60 (d, 1H), 12.35 (s, 1H) ppm.</p> <p>Melting point: 200-202°C.</p> <p>Crystallisation solvent: acetonitrile.</p>
13	-CH ₃		<p>Found: C, 53.40; H, 4.82; N, 15.37%. C₁₆H₁₇BrN₄O, requires C, 53.19; H, 4.74; N, 15.51%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.90 (s, 2H), 4.10 (s, 3H), 7.30 (m, 2H), 7.40 (d, 1H), 7.55 (s, 1H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 233-235 °C.</p> <p>Crystallisation solvent: ethanol.</p>

14	-CH ₃		<p>Found: C, 54.02; H, 5.15; N, 14.91%. C₁₇H₁₉BrN₄O, requires C, 54.41; H, 5.10; N, 14.93%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.75 (d, 3H), 1.90 (m, 2H), 2.95 (t, 2H), 4.10 (q, 1H), 4.25 (s, 3H), 7.23 (d, 2H), 7.50 (d, 2H), 9.50 (s, 1H) ppm.</p> <p>Melting point: 167-169°C.</p> <p>Crystallisation solvent: ethyl acetate/hexane.</p>
15	-CH ₃		<p>Found: C, 60.57; H, 5.39; N, 17.48%. C₁₆H₁₇ClN₄O, requires C, 60.66; H, 5.41; N, 17.69%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.90 (s, 2H), 4.07 (s, 3H), 7.35 (m, 4H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 189-191°C.</p> <p>Crystallisation solvent: acetonitrile.</p>

16	-CH ₃		<p>Found: C, 58.62; H, 4.98; N, 16.06%. C₁₇H₁₇F₃N₄O, requires C, 58.28; H, 4.89; N, 15.99%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.90 (t, 2H), 4.20 (s, 3H), 4.30 (s, 2H), 7.45 (m, 2H), 7.55 (m, 1H), 7.75 (d, 1H), 9.10 (s, 1H) ppm.</p> <p>Melting point: 195-196°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>
17	-CH ₃		<p>Found: C, 58.54; H, 4.91; N, 16.27%. C₁₇H₁₇F₃N₄O, requires C, 58.28; H, 4.89; N, 15.99%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.91 (t, 3H), 1.08 (m, 2H), 2.68 (t, 2H), 4.00 (s, 2H), 4.09 (s, 3H), 7.52 (d, 2H), 7.68 (d, 2H), 12.32 (s, 1H) ppm.</p> <p>Melting point: 209.5-211°C.</p> <p>Crystallisation solvent: ethyl acetate/hexane.</p>

18	-CH ₃		<p>Found: C, 70.06; H, 7.38; N, 17.28%. C₁₉H₂₄N₄O, requires C, 70.34; H, 7.46; N, 17.27%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.15 (d, 6H), 1.70 (m, 2H), 2.70 (m, 2H), 2.80 (m, 1H), 3.85 (s, 2H), 4.05 (s, 3H), 7.20 (m, 4H), 12.25 (s, 1H) ppm.</p> <p>Melting point: 209-210°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>
19	-CH ₃		<p>Found: C, 67.27; H, 7.10; N, 16.38%. C₁₉H₂₄N₄O₂, requires C, 67.03; H, 7.11; N, 16.38%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.25 (t, 3H), 1.85 (m, 2H), 2.95 (t, 2H), 2.55 (q, 2H), 4.05 (s, 2H), 4.25 (s, 3H), 4.50 (s, 2H), 7.35 (s, 4H), 9.80 (s, 1H) ppm.</p> <p>Melting point: 176-177 °C.</p>


20	-CH ₃		<p>Found: C, 66.42; H, 6.17; N, 17.11%. $C_{18}H_{20}N_4O_2$, requires C, 66.65; H, 6.22; N, 17.27%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.63 (s, 3H), 2.95 (t, 2H), 4.15 (s, 2H), 4.25 (s, 3H), 7.50 (d, 2H), 7.95 (d, 2H), 9.75 (s, 1H) ppm.</p> <p>Melting point: 209-210°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>
21	-CH ₃		<p>Found: C, 67.31; H, 7.52; N, 20.31%. $C_{19}H_{25}N_3O$, requires C, 67.23; H, 7.42; N, 20.63%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.25 (s, 6H), 2.95 (t, 2H), 3.65 (s, 2H), 4.05 (s, 2H), 4.25 (s, 3H), 7.30 (m, 4H), 9.30 (br.s, 1H) ppm.</p> <p>Melting point: 196-198°C.</p> <p>Crystallisation solvent: ethyl acetate.</p>

22	-CH ₃		<p>Found: C, 66.17; H, 7.17; N, 18.26%. C₂₁H₂₇N₃O₂, requires C, 66.12; H, 7.13; N, 18.36%. ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.45 (br, m, 4H), 2.95 (t, 2H), 3.50 (s, 2H), 3.75 (m, 4H), 4.10 (s, 2H), 4.25 (s, 3H), 7.35 (m, 4H), 9.35 (s, 1H) ppm. Melting point: 191-193°C. Crystallisation solvent: ethyl acetate/hexane.</p>
23	-CH ₃		<p>Found: C, 66.56; H, 6.24; N, 23.09%. C₂₀H₂₂N₂O, requires C, 66.28; H, 6.12; N, 23.19%. ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.85 (s, 2H), 4.05 (s, 3H), 5.15 (s, 2H), 6.85 (s, 1H), 7.05-7.30 (m, 5H), 7.70 (s, 1H), 12.25 (s, 1H) ppm. Melting point: 191-193°C. Crystallisation solvent: acetonitrile.</p>
24	-CH ₃		<p>Found: C, 47.27; H, 3.92; N, 17.16%. C₁₆H₁₆BrN₂O₃, requires C, 47.30; H, 3.97; N, 17.24%. ¹H-NMR (DMSO-d₆): δ = 0.80 (t, 3H), 2.55 (m, 2H), 2.55 (t, 2H), 4.05 (s, 3H), 4.30 (s, 2H), 7.50 (d, 1H), 7.95 (d, 1H), 8.25 (s, 1H), 12.35 (s, 1H) ppm. Melting point: 229-231°C. Crystallisation solvent: ethyl acetate.</p>

Example 25

5-(4-hydroxymethylbenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0097]



[0098] Aqueous hydrogen peroxide (0.24ml, 30% w/w, 0.0021mol) was added to a solution of sodium hydroxide (180mg, 0.0045mol) in water (10ml). A solution of N-(5-carbamoyl-1-methyl-3-propyl-1H-pyrazolyl)-2-(4-bromomethylphenyl)acetamide (400mg, 0.001mol) in dioxan (5ml) was then added and the mixture stirred at 90°C for 4 hours. On cooling, the reaction mixture was concentrated under reduced pressure, the residue dissolved in water (10ml) and acidified to pH 5 with 1N aqueous hydrochloric acid. This solution was then extracted with dichloromethane (2x20ml), the combined organic extracts dried (MgSO₄), filtered and evaporated under reduced pressure.

[0099] Purification by flash column chromatography, eluting with dichloromethane:methanol (96:4 by volume), followed by trituration with diethyl ether/pentane gave the title compound as a solid (60mg), m.p. 215-218°C.


[0100] Found: C, 65.14; H, 6.42; N, 17.88; C₁₇H₂₀N₄O₂, requires C, 65.36; H, 6.45; N, 17.94%.

[0101] ¹H-NMR (CDCl₃): δ = 1.07 (t, 3H), 1.86 (m, 3H), 2.92 (t, 2H), 4.08 (s, 2H), 4.24 (s, 3H), 4.70 (s, 2H), 7.34 (s, 4H), 9.74 (s, 1H) ppm.

Example 26

5-(2-methoxybenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0102]



[0103] N-(5-carbamoyl-1-methyl-3-propyl-1H-pyrazolyl)-2-(2-methoxyphenyl)acetamide (750mg, 0.00228mol) was suspended in polyphosphoric acid (10ml) and heated under nitrogen at 150°C for 4 hours. On cooling, the solution was added to ice and the pH adjusted to 5 with aqueous 10N sodium hydroxide solution. The aqueous phase was extracted with dichloromethane (3x10ml), the organic extracts combined, dried (MgSO₄), filtered and concentrated under reduced pressure.


[0104] Purification by flash column chromatography, eluting with dichloromethane:methanol (98:2, by volume), followed by trituration with hexane gave the title compound as a white solid, m.p. 162-163°C.

[0105] Found: C, 65.11; H, 6.41; N, 17.83; C₁₇H₂₀N₄O₂ requires C, 65.39; H, 6.45; N, 17.94%.

[0106] ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.90 (t, 2H), 4.00 (s, 3H), 4.05 (s, 2H), 4.20 (s, 3H), 7.00 (m, 2H), 7.35 (m, 2H), 9.50 (s, 1H) ppm.

Examples 27 to 29

[0107] The compounds of the following tabulated examples of the general formula: -



were prepared by reaction of the corresponding pyrazolocarboxamides by similar methods to that used in Example 26.

Ex. No.	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
27		<p>Found: C, 69.37; H, 7.17; N, 17.77%.</p> <p>C₁₈H₂₂N₄O, requires C, 69.65; H, 7.14; N, 18.05%.</p> <p>¹H-NMR (CDCl₃): δ = 0.95 (t, 3H), 1.05 (t, 3H), 1.90 (m, 2H), 2.05 (m, 1H), 2.40 (s, 1H), 2.95 (t, 2H), 3.80 (t, 1H), 4.30 (s, 3H), 7.20-7.45 (m, 5H), 10.15 (s, 1H) ppm.</p> <p>Melting point: 174-175°C.</p> <p>Crystallisation solvent: ethyl acetate/hexane.</p>
28		<p>Found: C, 70.10; H, 6.57; N, 18.30%.</p> <p>C₁₈H₂₀N₄O, requires C, 70.11; H, 6.54; N, 18.17%.</p> <p>¹H-NMR (CDCl₃): δ = 1.00 (t, 3H), 1.35 (m, 2H), 2.85 (m, 4H), 3.85 (t, 2H), 4.20 (s, 3H), 7.45 (m, 5H), 8.40 (s, 1H) ppm.</p> <p>Melting point: 122-124°C. Crystallisation solvent: ethanol.</p>
29		<p>Found: C, 69.41; H, 7.16; N, 17.65%.</p> <p>C₁₈H₂₂NO, requires C, 69.65; H, 7.14; N, 18.05%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.80 (s, 6H), 1.90 (m, 2H), 2.95 (t, 2H), 4.25 (s, 3H), 7.20-7.40 (m, 5H), 8.45 (s, 1H) ppm.</p> <p>Melting point: 175-178°C.</p>

Example 30

4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-benzoic acid

[0108]



[0109] N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-trifluoromethylphenyl)-acetamide (250mg, 0.00068mol) was suspended in polyphosphoric acid (4ml) and heated at 150°C for 4 hours under a nitrogen atmosphere. On cooling, the reaction mixture was neutralised with 1N aqueous sodium hydroxide solution (10ml) and extracted with dichloromethane/ methanol (40ml). The combined organic extracts were dried (MgSO₄), filtered and evaporated under reduced pressure. Crystallisation from methanol/ethyl acetate gave the title compound (166mg), m.p.292-293°C.


[0110] Found: C, 60.87; H, 5.50; N, 16.51; C₁₇H₁₆N₄O₃ · 0.5H₂O requires C, 60.89; H, 5.71; N, 16.71%.

[0111] ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.90 (m, 2H), 2.70 (t, 2H), 4.00 (s, 2H), 4.10 (s, 3H), 7.40 (d, 2H), 7.85 (d, 2H), 12.35 (s, 1H), 12.90 (br.s, 1H) ppm.

Example 31

5-(3-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0112]



[0113] 1-Methyl-5-(3-nitrobenzyl)-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (5.92g, 0.019mol) was suspended in industrial methylated spirit (300ml). Palladium on charcoal (10%, 610mg) was added and the resulting mixture was hydrogenated at 50 p.s.i and room temperature for 3 hours.


[0114] The resulting mixture was then filtered through Arbocel™, washing through with hot ethanol. The filtrate was concentrated under reduced pressure to give, after crystallisation from industrial methylated spirit, the title compound as a solid (5.53g), m.p.207-210°C.

[0115] Found: C, 64.54; H, 6.49; N, 23.09; C₁₆H₁₉N₅O, requires C, 64.62; H, 6.44; N, 23.55 %.

[0116] ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.70 (s, 2H), 4.05 (s, 3H), 5.00 (s, 2H), 6.40 (m, 3H), 6.90 (m, 1H), 12.20 (s, 1H) ppm.

Examples 32 and 33

[0117] The compounds of the following tabulated Examples of the general formula: -




were prepared from the corresponding nitroaromatic compounds by similar methods to that used in Example 31.

Ex. No.	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
32		<p>Found: C, 64.84; H, 6.54; N, 23.71%.</p> <p>C₁₆H₁₉N₅O, requires C, 64.62; H, 6.44; N, 23.55%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.70 (s, 2H), 4.05 (s, 3H), 5.30 (s, 2H), 6.50 (m, 1H), 6.65 (d, 1H), 6.95 (m, 1H), 7.10 (d, 1H), 12.20 (s, 1H) ppm.</p> <p>Melting point: 230-231°C. Crystallisation solvent: industrial methylated spirit.</p>
33		<p>Found: C, 63.40; H, 6.46; N, 22.92%.</p> <p>C₁₆H₁₉N₅O, requires C, 64.62; H, 6.44; N, 23.55%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.90 (t, 2H), 3.75 (s, 2H), 3.95 (s, 2H), 4.25 (s, 3H), 6.70 (d, 2H), 7.10 (d, 2H), 8.95 (s, 1H) ppm.</p> <p>Crystallisation solvent: acetone/hexane.</p>

Example 34

N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-phenyl}methanesulfonamide

[0118]



[0119] Methanesulfonyl chloride (43μl, 0.00055mol) was added to a solution of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (150mg, 0.0005mol) in pyridine (5ml), and the reaction stirred at room temperature, under a nitrogen atmosphere for 2 hours.


[0120] The reaction mixture was partitioned between saturated aqueous sodium carbonate solution (15ml) and dichloromethane (15ml), and the aqueous phase extracted with dichloromethane/methanol (50ml). The organic phases were combined, washed with 1N aqueous hydrochloric acid solution, dried (MgSO₄), filtered and concentrated under reduced pressure to give, after recrystallisation from ethyl acetate the title compound as a solid (81mg), m.p. 241-242°C.

[0121] Found: C, 54.34; H, 5.79; N, 18.84; C₁₇H₂₁N₅O₃S requires C, 54.38; H, 5.64; N, 18.65%.




[0122] ¹H-NMR (DMSO-d₆): δ = 0.95 (t, 3H), 1.75 (m, 2H), 2.80 (t, 2H), 2.85 (s, 3H), 3.90 (s, 2H), 4.15 (s, 3H), 7.20 (d, 2H), 7.30 (d, 2H), 9.00 (s, 1H), 11.15 (s, 1H) ppm.



Examples 35 to 45



[0123] The compounds of the following tabulated Examples of the general formula: -







were prepared by reaction of the appropriate anilines and sulfonyl or sulphamoyl chlorides using similar methods to that described in Example 34.

Ex. No.	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
35		<p>Found: C, 54.18; H, 5.61; N, 18.41%.</p> <p>C₁₇H₂₁N₃O₃S, requires C, 54.38; H, 5.64; N, 18.65%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 2.95 (s, 3H), 3.85 (s, 2H), 4.10 (s, 3H), 7.05 (m, 2H), 7.20 (s, 1H), 7.25 (m, 1H), 9.75 (s, 1H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 179-181°C. Crystallisation solvent: acetone/hexane.</p>
36		<p>Found: C, 54.90; H, 5.73; N, 18.79%.</p> <p>C₁₇H₂₁N₃O₃S, requires C, 54.38; H, 5.64; N, 18.65%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.05 (s, 3H), 4.05 (s, 2H), 4.10 (s, 3H), 7.15 (m, 1H), 7.25 (m, 2H), 7.40 (m, 1H) 9.80 (s, 1H), 12.40 (s, 1H) ppm.</p> <p>Melting point: 256-257°C. Crystallisation solvent: ethyl acetate.</p>
37		<p>Found: C, 60.31; H, 5.32; N, 15.96%.</p> <p>C₂₂H₂₃N₃O₃S, requires C, 60.40; H, 5.30; N, 16.01%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.75 (s, 2H), 4.05 (s, 3H), 7.00 (d, 2H), 7.15 (d, 2H), 7.55 (m, 3H), 7.75 (d, 2H), 10.20 (s, 1H), 12.20 (s, 1H) ppm.</p> <p>Melting point: 272-275°C. Crystallisation solvent: methanol/ethyl acetate.</p>

38		<p>Found: C, 60.67; H, 5.33; N, 16.04%.</p> <p>$C_{22}H_{23}N_5O_3S$, requires C, 60.40; H, 5.30; N, 16.01%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.75 (t, 2H), 3.70 (s, 2H), 4.10 (s, 3H), 7.00-7.25 (m, 4H), 7.50-7.80 (m, 5H), 10.30 (s, 1H), 12.35 (s, 1H) ppm</p> <p>Melting point: 244-248°C. Crystallisation solvent: methanol/ethyl acetate.</p>
39		<p>Found: C, 57.40; H, 6.51; N, 16.59%.</p> <p>$C_{20}H_{27}N_5O_3S$, requires C, 57.53; H, 6.52; N, 16.77%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.80 (t, 3H), 0.90 (t, 3H), 1.30 (m, 2H), 1.50-1.80 (m, 4H), 2.70 (t, 2H), 3.00 (t, 2H), 3.85 (s, 2H), 4.05 (s, 3H), 7.10 (d, 2H), 7.30 (d, 2H), 9.70 (s, 1H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 221-224°C. Crystallisation solvent: ethyl acetate/hexane.</p>

40		<p>Found: C, 52.48; H, 5.61; N, 15.91%.</p> <p>$C_{19}H_{24}N_5ClO_3S$, requires C, 52.11; H, 5.52; N, 15.99%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.10 (m, 2H), 2.75 (t, 2H), 3.15 (t, 2H), 3.70 (t, 2H), 3.85 (s, 2H), 4.10 (s, 3H), 7.10 (d, 2H), 7.30 (d, 2H), 9.80 (s, 1H), 12.25 (s, 1H) ppm.</p> <p>Melting point: 237-238°C. Crystallisation solvent: ethyl acetate.</p>
41		<p>Found: C, 52.41; H, 5.57; N, 15.76%.</p> <p>$C_{19}H_{24}N_5ClO_3S$, requires C, 52.11; H, 5.52; N, 15.99%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.15 (m, 2H), 2.70 (t, 2H), 3.25 (m, 2H), 3.70 (t, 2H), 4.05 (s, 2H), 4.10 (s, 3H), 7.15 (d, 1H), 7.25 (m, 2H), 7.40 (d, 1H), 9.90 (s, 1H), 12.40 (s, 1H) ppm.</p> <p>Melting point: 220-223°C. Crystallisation solvent: ethyl acetate.</p>


42		<p>Found: C, 57.15; H, 5.10; N, 19.12%.</p> <p>$C_{21}H_{22}N_2O_3S$, requires C, 57.52; H, 5.06; N, 19.16%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.80 (s, 2H), 4.05 (s, 3H), 7.00 (d, 2H), 7.20 (d, 2H), 7.55 (m, 1H), 8.10 (d, 1H), 8.75 (d, 1H), 8.85 (s, 1H), 10.45 (s, 1H), 12.20 (s, 1H) ppm.</p> <p>Melting point: 251-252°C. Crystallisation solvent: acetonitrile.</p>
43		<p>Found: C, 57.45; H, 5.04; N, 19.12%.</p> <p>$C_{21}H_{22}N_2O_3S$, requires C, 57.52; H, 5.06; N, 19.16%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.75 (s, 2H), 4.10 (s, 3H), 7.00 (d, 1H), 7.20 (m, 3H), 7.60 (m, 1H), 8.05 (d, 1H), 8.80 (m, 2H), 10.40 (s, 1H), 12.35 (s, 1H) ppm.</p> <p>Melting point: 233-234°C. Crystallisation solvent: ethyl acetate/hexane.</p>

44		<p>Found: C, 55.93; H, 6.58; N, 19.01%.</p> <p>$C_{20}H_{28}N_2O_3S$, requires C, 55.54; H, 6.52; N, 19.43%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.70 (t, 3H), 0.90 (t, 3H), 1.15 (m, 2H), 1.25 (m, 2H), 1.70 (m, 2H), 2.75 (m, 4H), 3.80 (s, 2H), 4.05 (s, 3H), 7.05 (d, 2H), 7.20 (d, 2H), 7.35 (m, 1H), 9.55 (s, 1H), 12.25 (s, 1H) ppm.</p> <p>Melting point: 174-183°C. Crystallisation solvent: ethyl acetate/hexane.</p>
45		<p>Found: C, 56.23; H, 6.24; N, 17.17%.</p> <p>$C_{19}H_{25}N_2O_3S$, requires C, 56.56; H, 6.25; N, 17.36%.</p> <p>1H-NMR (CDCl$_3$): δ = 1.05 (t, 3H), 1.20 (t, 3H), 1.85 (m, 2H), 2.90 (m, 5H), 3.75 (q, 2H), 4.05 (s, 2H), 4.25 (s, 3H), 7.35 (d, 2H), 7.45 (d, 2H), 9.50 (s, 1H) ppm.</p> <p>Melting point: 245-246°C. Crystallisation solvent: ethyl acetate.</p>

Example 46

N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}methanamide

[0124]



[0125] A mixture of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (297mg, 0.0010mol) and 1,2-diformylhydrazine (101mg, 0.00115mol) was heated to 200°C. The resulting melt was then stirred under a nitrogen atmosphere at this temperature for 55 minutes. On cooling, the solid was dissolved in dichloromethane: methanol, pre-absorbed onto silica and purified by flash column chromatography eluting with a solvent gradient of dichloromethane:methanol (99:1 to 96:4 by volume) then dichloromethane:methanol:0.880 aqueous ammonia (90:10:1 by volume). Crystallisation from ethyl acetate/methanol gave the title compound (170mg), m.p. 260-261°C.

[0126] Found: C, 62.44; H, 5.93; N, 21.22; $C_{17}H_{19}N_5O_2$, requires C, 62.76; H, 5.89; N, 21.52%


[0127] 1H -NMR (DMSO- d_6): δ = 0.91 (t, 3H), 1.70 (m, 2H), 2.82 (t, 2H), 3.84 (s, 2H), 4.06 (s, 3H), 7.25 (d, 2H), 7.50

(d, 2H), 8.22 (s, 1H), 10.16 (s, 1H), 12.28 (s, 1H) ppm.

Example 47

N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}acetamide

[0128]



[0129] Triethylamine (160 μ l, 0.0011mol), dimethylaminopyridine (10mg), and acetic anhydride (50 μ l, 0.0005mol) were added to a solution of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (110mg, 0.00037mol) in dichloromethane (9ml). The resulting mixture was stirred overnight at room temperature under a nitrogen atmosphere. The mixture was then partitioned between 2M aqueous hydrochloric acid solution (15ml) and dichloromethane (15ml) and the aqueous phase extracted with dichloromethane/methanol (30ml). The combined organic extracts were dried (MgSO₄), filtered and concentrated under reduced pressure to give, after crystallisation from ethyl acetate/methanol, the title compound as an off white solid (105mg), m.p.272-275°C.


[0130] Found: C, 63.17; H, 6.25; N, 20.05; C₁₈H₂₁N₅O₂, requires C, 63.70; H, 6.24; N, 20.64 %.

[0131] ¹H-NMR (DMSO-d₆): δ = 0.92 (t, 3H), 1.70 (t, 2H), 2.00 (s, 3H), 2.72 (t, 2H), 3.82 (s, 2H), 4.06 (s, 3H), 7.21 (d, 2H), 7.50 (d, 2H), 9.88 (s, 1H), 12.24 (s, 1H) ppm.

Example 48

5-[4-(ethylamino)benzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0132]



[0133] N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}acetamide (2.37g, 0.007mol) was added to a mixture of lithium aluminium hydride (705mg, 0.0186mol) in tetrahydrofuran (250ml), under a nitrogen atmosphere. After the effervescence had subsided, the reaction was refluxed for 2 hours. On cooling, the mixture was treated dropwise with aqueous sodium hydroxide solution (5M, 1.1ml), and the resulting solid filtered and washed well with tetrahydrofuran (250ml). The filtrate was concentrated under reduced pressure and the residue partitioned between ethyl acetate (100ml) and water (100ml). The aqueous phase was extracted with ethyl acetate (200ml) and the organic phases combined, dried (MgSO₄), filtered and concentrated under reduced pressure.

[0134] Purification by flash column chromatography, eluting with ethyl acetate followed by crystallisation from ethyl acetate gave the title compound (1.03g), m.p.186-188°C.

[0135] Found: C, 66.57; H, 7.09; N, 21.54; C₁₈H₂₃N₅O, requires C, 66.44; H, 7.12; N, 21.52%.

[0136] ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.10 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 2.95 (m, 2H), 3.70 (s, 2H), 4.05

(s, 3H), 5.40 (t, 1H), 6.45 (d, 2H), 7.00 (d, 2H), 12.15 (s, 1H) ppm.

Example 49

N-ethyl-N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}acetamide

[0137] The title compound was prepared from 5-[4-(ethylamino)benzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo [4,3-d]pyrimidin-7-one following the procedure described in Example 47 and was obtained, after trituration with ethyl acetate as a solid (59%), m.p. 149-151°C.


[0138] Found: C, 65.05; H, 6.88; N, 18.92; $C_{20}H_{25}N_5O_2$, requires C, 65.37; H, 6.86; N, 19.06%.

[0139] 1H -NMR ($CDCl_3$): δ = 1.05 (t, 3H), 1.15 (t, 3H), 1.86 (m, 5H), 2.92 (t, 2H), 3.76 (q, 2H), 7.18 (d, 2H), 7.44 (d, 2H), 9.75 (s, 1H) ppm.

Example 50

5-[4-(diethylamino)benzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0140]



[0141] The title compound was prepared from N-ethyl-N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo [4,3-d]pyrimidin-5-yl)methyl]phenyl}acetamide following the procedure described in Example 48 and was obtained after crystallisation from acetone/hexane as a solid (45%), m.p. 201-203°C.


[0142] Found: C, 68.11; H, 7.72; N, 19.94; $C_{20}H_{27}N_5O$ requires C, 67.96; H, 7.70; N, 19.82%.

[0143] 1H -NMR ($CDCl_3$): δ = 1.05 (t, 3H), 1.20 (t, 6H), 1.85 (m, 2H), 2.90 (t, 2H), 3.40 (q, 4H), 3.95 (s, 2H), 4.20 (s, 3H), 6.70 (d, 2H), 7.10 (d, 2H), 8.65 (s, 1H) ppm.

Example 51

N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-2-nitrophenyl}acetamide

[0144]



[0145] Glacial acetic acid (1.93ml, 0.0337mol) was added to N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}acetamide (1.93g, 0.0057 mol), followed by concentrated sulphuric acid (5.8ml) and the mixture cooled in an icebath. Concentrated nitric acid (3.86ml) was then added dropwise and once addition was complete, the reaction was stirred at room temperature for an hour.

[0146] The mixture was carefully poured onto ice and then extracted with dichloromethane (2x50ml). The combined organic extracts were dried (MgSO_4), filtered and evaporated under reduced pressure.

[0147] Purification by flash column chromatography eluting with methanol:dichloromethane (2:98 by volume), followed by recrystallisation from acetic acid/water gave the title compound (619mg), m.p. 261-262°C.


[0148] Found: C, 55.84; H, 5.17; N, 21.48; $\text{C}_{18}\text{H}_{20}\text{N}_6\text{O}_4$, requires C, 56.24; H, 5.24; N, 21.87%

[0149] $^1\text{H-NMR}$ (DMSO-d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.05 (s, 3H), 2.71 (t, 2H), 3.99 (s, 2H), 4.08 (s, 3H), 7.53 (d, 1H), 7.60 (d, 1H), 10.23 (s, 1H) ppm.

Example 52

2-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenylamino}acetic acid.

[0150]



[0151] 10% w/w Palladium on charcoal (87mg) was added to a solution of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (870mg, 0.00293mol) and glyoxylic acid hydrate (260mg, 0.0035mol) in methanol (20ml) and the mixture hydrogenated at room temperature at 50 p.s.i. for 3 hours.

[0152] The mixture was filtered through Arbocel™ and washed well with 10% aqueous sodium hydroxide (60ml). The filtrate was evaporated under reduced pressure and the residue partitioned between water (40ml) and dichloromethane (40ml). The aqueous layer was acidified to pH1 with 2M aqueous hydrochloric acid and extracted well with dichloromethane (2x50ml). These combined organic extracts were dried (MgSO_4), filtered and evaporated under reduced pressure to give a brown solid.


[0153] Purification by flash column chromatography eluting with methanol:dichloromethane (3:97 by volume), followed by recrystallisation from acetone/hexane gave the title compound (180mg), m.p. 175-178°C.

[0154] $^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (t, 3H), 1.87 (m, 2H), 2.92 (t, 2H), 3.82 (s, 2H), 3.96 (m, 3H), 4.24 (s, 3H), 6.64 (d, 2H), 7.14 (d, 2H), 8.67 (s, 1H) ppm.

Example 53

N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}-(E)-3-ethoxy-2-propenamide

[0155]



[0156] (E)-3-ethoxyacryloyl chloride (*J. Chem. Soc.*; 1958,153) (298mg, 0.0022mol), was added dropwise to an ice-cooled solution of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (600mg, 0.002mol) in pyridine (15ml) and the reaction stirred at room temperature for 20 hours. The mixture was partitioned between water (30ml) and dichloromethane (30ml), the aqueous layer acidified to pH1 with 2M aqueous hydrochloric acid solution and then extracted with dichloromethane (2x40ml). The combined organic extracts were dried (MgSO_4), filtered and evaporated under reduced pressure, to give, after recrystallisation from ethyl acetate/methanol the title compound (483mg), m.p. 241-243°C


[0157] Found: C, 63.92; H, 6.44; N, 17.61; $\text{C}_{21}\text{H}_{12}\text{N}_5\text{O}_3$, requires C, 63.78; H, 6.37; N, 17.71%

[0158] $^1\text{H-NMR}$ (DMSO-d_6): δ = 0.93 (t, 3H), 1.24 (t, 3H), 1.70 (m, 2H), 2.74 (t, 2H), 3.82 (s, 2H), 3.94 (q, 2H), 4.08 (s, 3H), 5.50 (d, 1H), 7.21 (d, 2H), 7.44 (d, 1H), 7.52 (d, 2H), 9.68 (s, 1H), 12.24 (s, 1H) ppm.

Example 54

1-methyl-5-[(2-oxo-1,2-dihydro-6-quinolinyl)methyl]-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0159]



[0160] Concentrated sulphuric acid (10ml) was added to N-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}-(E)-3-ethoxy-2-propenamide (980mg, 0.00248mol) and the reaction stirred at room temperature for 4 hours. The mixture was then poured carefully onto ice and basified with 0.880 aqueous ammonia solution. The resulting precipitate was filtered and then triturated with boiling acetic acid/water to give the title compound (460mg), m.p. > 300°C


[0161] Found: C, 64.57; H, 5.52; N, 19.94; $\text{C}_{19}\text{H}_{19}\text{N}_5\text{O}_2 \cdot 0.25\text{H}_2\text{O}$, requires C, 64.47; H, 5.55; N, 19.79%

[0162] $^1\text{H-NMR}$ (DMSO-d_6): δ = 0.91 (t, 3H), 1.68 (m, 2H), 2.70 (t, 2H), 3.94 (s, 2H), 4.09 (s, 3H), 6.44 (d, 1H), 7.22 (d, 1H), 7.45 (d, 1H), 7.56 (s, 1H), 7.84 (d, 1H), 11.70 (s, 1H), 12.22 (s, 1H) ppm.

Example 55

N-{2-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-phenyl}-3-(dimethylamino)-1-propanesulfonamide

[0163]



[0164] N-{2-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-phenyl}-3-chloro-1-propanesulfonyl chloride (200mg, 0.00046mol) was dissolved in 33% w/w ethanolic dimethylamine (10ml) and the reaction stirred at reflux for 3 hours. On cooling, the reaction was concentrated under reduced pressure and the residue partitioned between water (15ml) and dichloromethane (15ml). The aqueous layer was extracted with dichloromethane (40ml) and the combined organic extracts dried (MgSO_4), filtered and evaporated under reduced pressure.

[0165] Purification by flash column chromatography eluting with a solvent gradient of 0.880 aqueous ammonia:methanol:dichloromethane (0:3:97 to 1:10:89 by volume) gave, after recrystallisation from ethyl acetate/hexane, the title compound (70mg), m.p. 202-204°C.


[0166] Found: C, 56.08; H, 6.80; N, 18.74; $\text{C}_{21}\text{H}_{30}\text{N}_6\text{O}_3\text{S}$, requires C, 56.48; H, 6.77; N, 18.82%

[0167] $^1\text{H-NMR}$ (CDCl_3): δ = 1.08 (t, 3H), 1.86 (m, 2H), 2.12 (m, 2H), 2.23 (s, 6H), 2.46 (t, 2H), 2.94 (t, 2H), 3.29 (t, 2H), 4.16 (s, 2H), 4.32 (s, 3H), 7.10 (m, 1H), 7.28 (m, 1H), 7.48 (d, 1H), 7.57 (d, 1H) ppm.

Example 56

2-{2-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-phenyl}tetrahydro-2H-isothiazole-1,1-dioxide

[0168]



[0169] N-{2-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-phenyl}-3-chloro-1-propanesulfonyl chloride (200mg, 0.00046mol) was added to a mixture of sodium hydride (27mg, 80% w/w oil dispersion, 0.00091mol) in dimethylformamide (10ml) and the reaction stirred for 20 hours at room temperature, under a nitrogen atmosphere. Methanol (5ml) was added and the mixture concentrated under reduced pressure. The residue was partitioned between ethyl acetate (20ml) and water (20ml), the aqueous phase acidified to pH1 with aqueous hydrochloric acid solution (1M) and extracted with ethyl acetate (2x25ml). These combined organic phases were dried (MgSO_4), filtered, and concentrated under reduced pressure to give, after recrystallisation from ethyl acetate/hexane the title compound, as a white solid (139mg), m.p. 181-182°C.

[0170] Found: C, 56.40; H, 5.77; N, 17.34; $C_{19}H_{23}N_5O_3S$, requires C, 56.84; H, 5.77; N, 17.44%.

[0171] 1H -NMR (DMSO- d_6): δ = 0.85 (t, 3H), 1.65 (m, 2H), 2.35 (m, 2H), 2.65 (t, 2H), 3.40 (t, 2H), 3.60 (t, 2H), 4.05 (s, 2H), 4.10 (s, 3H), 7.35 (m, 3H), 7.45 (m, 1H), 12.20 (s, 1H) ppm.

5 Example 57

2-{4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl}tetrahydro-2H-isothiazole-1,1-dioxide

[0172] The title compound was prepared from N-(4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl)-3-chloro-1-propanesulfonyl chloride following the procedure described in Example 56 and was obtained as a solid (70%), m.p. 254-255°C.


[0173] Found: C, 56.73; H, 5.77; N, 17.30; $C_{19}H_{23}N_5O_3S$, requires C, 56.84; H, 5.77; N, 17.44%.

[0174] 1H -NMR (DMSO- d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.35 (m, 2H), 2.70 (t, 2H), 3.45 (t, 2H), 3.70 (t, 2H), 3.85 (s, 2H), 4.05 (s, 3H), 7.15 (d, 2H), 7.35 (d, 2H), 12.30 (s, 1H) ppm.


Examples 58 and 59

Preparation of 3-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-benzenesulfonyl chloride and 4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-benzenesulfonyl chloride

[0175]



and



[0176] 5-benzyl-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (500 mg, 0.00178 mol) was dissolved in chlorosulfonic acid (2.5 ml, 0.0376 mol) to give an orange solution which was then warmed to 60°C for 2 hours. On cooling, the solution was pipetted onto ice and then extracted with dichloromethane (2x20 ml). The combined organic extracts were dried ($MgSO_4$), filtered and concentrated under reduced pressure. Crystallisation from ethyl acetate gave the 4-substituted title compound, m.p. >350°C (some decomposition at 240°C).

[0177] Found: C, 50.56; H, 4.51; N, 14.48; $C_{16}H_{17}N_4ClO_3S$, requires C, 50.46; H, 4.50; N, 14.71%.

[0178] 1H -NMR ($CDCl_3$): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.90 (t, 2H), 4.10 (s, 2H), 4.30 (s, 3H), 7.70 (d, 2H), 8.00 (d, 2H), 11.60 (s, 1H) ppm.

[0179] The mother liquors were evaporated under reduced pressure to give the 3-substituted title compound (88 mg).


[0180] Found: C, 50.56; H, 4.51; N, 14.48; $C_{16}H_{17}N_4ClO_3S$, requires C, 50.46; H, 4.50; N, 14.71%.

[0181] $^1\text{H-NMR}$ (CDCl_3): $\delta = 1.03$ (t, 3H), 1.86 (m, 2H), 2.92 (t, 2H), 4.22 (s, 2H), 4.33 (s, 3H), 7.60 (m, 1H), 7.86 (d, 1H), 8.00 (d, 1H), 8.23 (s, 1H), 11.98 (s, 1H) ppm.


Examples 60 and 61

3-[(1-ethyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-1-benzenesulfonamide and 4-[(1-ethyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-1-benzenesulfonamide

[0182]



and



[0183] 5-benzyl-1-ethyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (480mg, 0.0016 mol) was dissolved in chlorosulfonic acid (5ml) and the solution warmed to 60°C for 2 hours. On cooling, the mixture was pipetted onto ice to give a pale brown suspension.

[0184] 0.880 Aqueous ammonia solution was then added until the mixture became basic with dissolution of all the solid. The resulting orange solution was then acidified with concentrated hydrochloric acid solution and extracted with dichloromethane/methanol (2x75ml). The organic extracts were combined, dried (MgSO_4), filtered and concentrated under reduced pressure.

[0185] Purification by flash column chromatography, eluting with a solvent gradient of dichloromethane/methanol (97:3 to 95:5, by volume), gave a fractional separation of the two product isomers. The first product fractions to elute were combined, evaporated under reduced pressure, the residue was crystallised from ethyl acetate to give the 3-substituted title compound as a solid (40mg), m.p. 222-224°C.

[0186] Found: C, 54.20; H, 5.68; N, 18.57; $\text{C}_{17}\text{H}_{21}\text{N}_5\text{O}_3\text{S}$, requires C, 54.39; H, 5.64; N, 18.65%.

[0187] $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): $\delta = 0.90$ (t, 3H), 1.35 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 4.00 (s, 2H), 4.45 (m, 2H), 7.30 (s, 2H), 7.50 (m, 2H), 7.70 (d, 1H), 7.80 (s, 1H), 12.40 (s, 1H) ppm.

[0188] The later product fractions to elute were combined, evaporated under reduced pressure, and the residue was crystallised from methanol, to give the 4-substituted title compound as a solid (110mg), m.p. 248-250°C.


[0189] Found: C, 54.26; H, 5.63; N, 18.52; $\text{C}_{17}\text{H}_{21}\text{N}_5\text{O}_3\text{S}$, requires C, 54.39; H, 5.64; N, 18.65%.

[0190] $^1\text{H-NMR}$ ($\text{DMSO}-d_6$): $\delta = 0.90$ (t, 3H), 1.30 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 4.00 (s, 2H), 4.45 (m, 2H), 7.30 (s, 2H), 7.50 (d, 2H), 7.75 (d, 2H), 12.35 (s, 1H) ppm.

Example 62

5-(4-([4-methylpiperazin-1-yl]sulphonyl)benzyl)-1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidine

[0191]



[0192] N-methyl piperazine (1.28ml, 0.0115mol) was added to a solution of 4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-1-benzenesulfonyl chloride (1.35g, 0.0035mol) in ethanol (20ml) and the reaction stirred at room temperature for 18 hours. The reaction mixture was concentrated under reduced pressure and the residue partitioned between dichloromethane (50ml) and water (20ml). The aqueous phase was extracted with further dichloromethane (50ml), the combined organic extracts dried (MgSO₄), filtered, and evaporated under reduced pressure.


[0193] Purification by flash column chromatography eluting with methanol:dichloromethane (3:97 by volume) followed by trituration with ethyl acetate gave the title compound as a solid.

[0194] Found: C, 55.91; H, 6.29; N, 18.53; C₂₁H₂₈N₆O₃S, requires C, 56.74; H, 6.35; N, 18.90%.



[0195] ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.25 (s, 3H), 2.45 (m, 4H), 3.00 (m, 6H), 4.15 (s, 2H), 4.30 (s, 3H), 7.68 (m, 4H), 11.60 (s, 1H) ppm.



Examples 63 to 66

[0196] The compounds of the following tabulated Examples of the general formula:-



were prepared by reaction of the corresponding sulfonyl chloride and amine by similar methods to that used in Example 62.


Ex. No.	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
63		<p>Found: C, 59.10; H, 5.59; N, 17.91%.</p> <p>C₂₃H₂₆N₆O₃S, requires C, 59.21; H, 5.62; N, 18.01%.</p> <p>¹H-NMR (CDCl₃): δ = 1.00 (t, 3H), 1.70 (m, 2H), 2.65 (s, 3H), 2.90 (t, 2H), 4.15 (s, 2H), 4.25 (s, 2H), 4.30 (s, 3H), 7.20 (m, 1H), 7.50 (m, 2H), 7.75 (m, 3H), 8.00 (s, 1H), 8.40 (s, 1H), 11.60 (s, 1H) ppm.</p> <p>Melting point: 180-182°C. Crystallisation solvent: ethyl acetate/methanol.</p>
64		<p>Found: C, 58.96; H, 5.63; N, 17.90%.</p> <p>C₂₃H₂₆N₆O₃S, requires C, 59.21; H, 5.62; N, 18.01%.</p> <p>¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.70 (s, 3H), 2.95 (t, 2H), 4.15 (s, 2H), 4.30 (s, 5H), 7.20 (m, 1H), 7.55 (d, 1H), 7.65 (d, 2H), 7.73 (m, 1H), 7.82 (d, 2H), 8.47 (s, 1H), 11.40 (s, 1H) ppm.</p> <p>Melting point: 206-208°C. Crystallisation solvent: ethyl acetate/methanol.</p>

65		<p>Found: C, 50.72; H, 5.51; N, 14.67%.</p> <p>$C_{20}H_{25}N_3O_6S \cdot 0.5 H_2O$, requires C, 50.83; H, 5.55; N, 14.82%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.28 (s, 3H), 3.45 (m, 2H), 3.80 (m, 1H), 4.00 (s, 2H), 4.10 (s, 3H), 5.05 (m, 1H), 7.50 (d, 2H), 7.70 (d, 2H), 8.20 (br.s, 1H), 12.35 (br.s, 1H) ppm.</p> <p>Melting point: 225-227°C. Crystallisation solvent: ethyl acetate/methanol.</p>
66		<p>Found: C, 51.44; H, 5.49; N, 14.74%.</p> <p>$C_{20}H_{25}N_3O_6S$, requires C, 51.83; H, 5.44; N, 15.11%.</p> <p>1H-NMR (DMSO-d_6): δ = 0.90 (t, 3H), 1.65 (m, 2H), 2.70 (t, 2H), 3.30 (s, 3H), 3.45 (m, 2H), 3.80 (t, 1H), 4.00 (s, 2H), 4.10 (s, 3H), 5.00 (m, 1H), 7.50 (m, 2H), 7.65 (m, 1H), 7.75 (s, 1H), 8.25 (br.s, 1H), 12.40 (br.s, 1H) ppm.</p> <p>Melting point: 148-150°C. Crystallisation solvent: ethyl acetate/methanol.</p>

Example 67

4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-1-benzenesulfonic acid

[0197]



[0198] A solution of 4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]-1-benzenesulfonyl chloride (250mg, 0.00065mol) in 1N aqueous sodium hydroxide solution (3ml) was stirred at room temperature for 2 hours. The reaction was acidified to pH1 with 2N aqueous hydrochloric acid and the resulting crystals filtered, washed with water, and dried by air suction, to yield the title compound (178mg), m.p. > 350°C.


[0199] Found: C, 52.35; H, 5.04; N, 15.13; $C_{16}H_{18}N_4O_4S$, requires: C, 53.03; H, 5.01; N, 15.46%

[0200] 1H -NMR (DMSO- d_6): δ = 0.90 (t, 3H), 1.68 (m, 2H), 2.71 (t, 2H), 3.92 (s, 2H), 4.07 (s, 3H), 5.80 (s, 1H), 7.25 (d, 2H), 7.54 (d, 2H) ppm.

Example 68

(2S)-3-hydroxy-2-((4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl)sulfonamido)propanamide

[0201]



[0202] Methyl (2S)-3-hydroxy-2-((4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl)sulfonamido)propanoate (280mg, 0.0006mol) was dissolved in 0.880 aqueous ammonia solution (10ml), and stirred at room temperature for 60 hours. The reaction mixture was then concentrated under reduced pressure, suspended in water (10ml), then re-concentrated under reduced pressure. Crystallisation from ethanol gave the title compound (192mg), m.p. 226-229°C.

[0203] Found: C, 49.97; H, 5.70; N, 18.04; $C_{19}H_{24}N_6O_5S \cdot 0.5 H_2O$, requires C, 49.87; H, 5.51; N, 18.37%.

[0204] 1H -NMR (DMSO- d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.40 (m, 2H), 3.60 (m, 1H), 3.95 (s, 2H), 4.10 (s, 3H), 4.85 (m, 1H), 7.00 (s, 1H), 7.20 (s, 1H), 7.45 (d, 2H), 7.65 (d, 1H), 7.75 (d, 2H), 12.35 (s, 1H) ppm.

Example 69

(2S)-3-hydroxy-2-((3-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl)sulfonamido)propanamide

[0205] The title compound was prepared from methyl-(2S)-3-hydroxy-2-((3-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenyl)sulfonamido)propanoate following the procedure described in Example 68 and was obtained as a solid (40%) m.p. 232-238°C.


[0206] Found: C, 49.24; H, 5.73; N, 17.82; $C_{19}H_{24}N_6O_5S \cdot H_2O$, requires C, 48.92; H, 5.62; N, 18.01%.

[0207] 1H -NMR (DMSO- d_6): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 3.40 (m, 2H), 3.65 (m, 1H), 4.00 (s, 2H), 4.10 (s, 3H), 4.85 (m, 1H), 7.05 (s, 1H), 7.20 (s, 1H), 7.50 (m, 2H), 7.70 (m, 2H), 7.80 (s, 1H), 12.40 (s, 1H) ppm.

Example 70

5-[4-(4H-1,2,4-triazol-4-yl)benzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0208]



[0209] A mixture of 5-(4-aminobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (297mg, 0.0010mol) and 1,2-diformylhydrazine (101mg, 0.00115mol) was heated to 200°C. The resulting melt was then stirred under a nitrogen atmosphere at this temperature for 55 minutes. On cooling, the solid was dissolved in dichloromethane: methanol, pre-absorbed onto silica and purified by flash column chromatography eluting with a solvent gradient of dichloromethane:methanol (99:1 to 96:4 by volume) then dichloromethane:methanol:0.880 aqueous ammonia (90:10:1 by volume).

[0210] Crystallisation from ethyl acetate/methanol gave the title compound (120mg), m.p.297-300°C.


[0211] Found: C, 60.87; H, 5.56; N, 27.21; C₁₈H₁₉N₇O 0.25H₂O requires C, 61.09; H, 5.55; N, 27.70%.

[0212] ¹H-NMR (DMSO-d₆): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.72 (t, 2H), 3.96 (s, 2H), 4.08 (s, 3H), 7.50 (d, 2H), 7.63 (d, 2H), 9.07 (s, 2H), 12.35 (s, 1H).

Example 71

5-[4-(1-Imidazolyl)benzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0213]



[0214] 5-(4-Bromobenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (390mg, 0.00108mol), imidazole (380mg, 0.00558mol), potassium carbonate (160mg, 0.00116mol), copper bronze (75mg) and iodine (46mg, 0.00018mol) in N-methyl-2-pyrrolidinone (6ml) were heated under a nitrogen atmosphere at 200°C for 3 hours. On cooling, the mixture was concentrated under reduced pressure and the residue partitioned between ethyl acetate (10ml) and water (10ml). The aqueous phase was extracted with ethyl acetate (30ml), and the combined organic extracts washed with brine, dried (MgSO₄), filtered and concentrated under reduced pressure.


[0215] Purification by flash column chromatography eluting with 0.880 aqueous ammonia:methanol:dichloromethane (0.5:5:95, by volume) followed by crystallisation from ethanol gave the title compound as a solid (85mg) m.p.252-254°C.

[0216] Found: C, 65.28; H, 5.76; N, 24.01; C₁₉H₂₀N₆O, requires C, 65.50; H, 5.79; N, 24.12%.

[0217] ¹H-NMR (CDCl₃): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.95 (t, 2H), 4.15 (s, 2H), 4.30 (s, 3H), 7.30 (m, 2H), 7.40 (d, 2H), 7.50 (d, 2H), 7.83 (s, 1H), 10.45 (s, 1H) ppm.



Examples 72 to 74

[0218] The compounds of the following tabulated examples of the general formula:-



were prepared by reaction of the appropriate bromides and heterocycles following the procedure described in Example 71.


Ex. No.	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting Point/Crystallisation solvent
72		<p>Found: C, 62.14; H, 5.53; N, 28.30%.</p> <p>C₁₈H₁₉N₇O, requires C, 61.88; H, 5.48; N, 28.06%.</p> <p>¹H-NMR (CDCl₃): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.70 (t, 2H), 4.00 (s, 2H), 4.10 (s, 3H), 7.50 (d, 2H), 7.80 (d, 2H), 8.20 (s, 1H), 9.20 (s, 1H), 12.35 (s, 1H) ppm.</p> <p>Melting point: 221-224°C. Crystallisation solvent: ethyl acetate/methanol.</p>

73		<p>Found: C, 66.26; H, 6.26; N, 23.14%.</p> <p>$C_{20}H_{22}N_6O$, requires C, 66.28; H, 6.12; N, 23.19%.</p> <p>1H-NMR ($CDCl_3$): δ = 1.05 (t, 3H), 1.80 (d, 3H), 1.90 (m, 2H), 2.95 (t, 2H), 4.20 (q, 1H), 4.25 (s, 3H), 7.30 (m, 2H), 7.35 (d, 2H), 7.50 (d, 2H), 7.80 (s, 1H), 10.20 (s, 1H) ppm.</p> <p>Melting point: 210-212°C.</p>
74		<p>Found: C, 65.98; H, 6.13; N, 22.95%</p> <p>$C_{20}H_{22}N_6O$, requires C, 66.28; H, 6.12; N, 23.19%</p> <p>1H-NMR ($DMSO-d_6$): δ = 0.90 (t, 3H), 1.70 (m, 2H), 2.15 (s, 3H), 2.70 (t, 2H), 3.90 (s, 2H), 4.10 (s, 3H), 7.40 (m, 3H), 7.55 (d, 2H), 8.05 (s, 1H), 12.30 (s, 1H) ppm.</p> <p>Melting point: 244-246°C. Crystallisation solvent: ethyl acetate/methanol.</p> <p>(Only the 4-isomer was isolated)</p>

Example 75

5-(4-hydroxybenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0219]



[0220] 60% w/w Hydrobromic acid (20ml) was added to 5-(4-methoxybenzyl)-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (630mg) and the mixture stirred at 130°C for 90 minutes. On cooling, the reaction was neutralized with saturated aqueous sodium carbonate solution and partitioned between dichloromethane (40ml) and water (40ml). The aqueous layer was further extracted with dichloromethane (100ml), the combined organic layers dried ($MgSO_4$), filtered and evaporated under reduced pressure. Crystallisation from ethyl acetate gave the title compound (75mg), m.p. 260°C.


[0221] Found: C, 63.35; H, 6.02; N, 18.68; $C_{16}H_{18}N_4O_2 \cdot 0.25H_2O$, requires: C, 63.46; H, 6.16; N, 18.50%.

[0222] 1H -NMR ($CDCl_3$): δ = 0.95 (t, 3H), 1.74 (m, 2H), 2.80 (t, 2H), 3.83 (s, 2H), 4.12 (s, 3H), 6.72 (d, 2H), 7.08 (d, 2H), 8.60 (s, 1H), 10.70 (s, 1H) ppm.

Example 76

4-[(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)methyl]phenoxyacetic acid

[0223]



[0224] Aqueous hydrogen peroxide solution (0.35ml, 30% w/w, 0.0031mol) was added to a solution of sodium hydroxide (350mg, 0.0088mol) in water (14ml). Tert-butyl 2-(4-{N-[5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl]carbamoyl}methyl)phenoxy)acetate (530mg, 0.0012mol) in ethanol (7ml) was then added and the reaction stirred at reflux for 1 hour. On cooling, the mixture was concentrated under reduced pressure, the residue dissolved in water (15ml) and acidified to pH2 with 1N aqueous hydrochloric acid.

[0225] The resulting precipitate was filtered, washed with water (40ml) and dried under air suction. Recrystallisation from ethanol gave the title compound (340mg), m.p.230-232°C.


[0226] Found: C, 57.69; H, 5.52; N, 14.90; $C_{18}H_{20}N_4O_4 \cdot H_2O$, requires C, 57.75; H, 5.92; N, 14.97%.

[0227] 1H -NMR (DMSO- d_6): δ = 0.92 (t, 3H), 1.68 (m, 2H), 2.72 (t, 2H), 3.81 (s, 2H), 4.07 (s, 3H), 4.62 (s, 2H), 6.82 (d, 2H), 7.23 (d, 2H), 12.25 (s, 1H), 12.98 (s, 1H) ppm.

Example 77

(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methyl acetate

[0228]



[0229] 2-chloro-2-phenylacetyl chloride (3.47ml, 0.022mol) was added dropwise to a solution of 4-amino-3-propyl-1-methyl-5-pyrazolecarboxamide (4g, 0.022mol) in acetic acid (30ml) and the reaction stirred at reflux under a nitrogen atmosphere for 20 hours.

[0230] On cooling, the mixture was concentrated under reduced pressure and purified by flash column chromatography eluting with dichloromethane. Trituration with diethyl ether gave the title compound (633mg), m.p.161-163°C.


[0231] Found: C, 63.37; H, 5.84; N, 16.07; $C_{18}H_{20}N_4O_3$, requires C, 63.51; H, 5.92; N, 16.46%

[0232] 1H -NMR (CDCl $_3$): δ = 1.01 (t, 3H), 1.82 (m, 2H), 2.30 (s, 3H), 2.90 (t, 2H), 4.24 (s, 3H), 6.74 (s, 1H), 7.40 (m, 3H), 7.50 (m, 2H), 9.88 (s, 1H) ppm.

Example 78

5-[α -hydroxybenzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one

[0233]



[0234] (1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)methyl acetate (1.311g, 0.004mol) was added to a solution of potassium hydroxide (232mg, 0.0041mol) in ethanol (60ml). The resulting mixture was refluxed for 1 hour to give a colourless solution. On cooling, this was concentrated under reduced pressure, and the resulting residue partitioned between water and dichloromethane. The aqueous phase was acidified to pH3 with 1M aqueous hydrochloric acid solution and was extracted with dichloromethane/methanol (100ml). The organic extracts were combined, dried (MgSO_4), filtered and concentrated under reduced pressure.

[0235] Purification by flash column chromatography eluting with dichloromethane:methanol (95:5 by volume), followed by crystallisation from acetone/hexane, gave the title compound (50mg), m.p.164-165°C.


[0236] Found: C, 64.29; H, 6.10; N, 19.24; $\text{C}_{16}\text{H}_{16}\text{N}_4\text{O}_2$ requires C, 64.41; H, 6.08; N, 18.78%.

[0237] $^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (t, 3H), 1.85 (m, 2H), 2.90 (t, 2H), 4.05 (s, 1H), 4.25 (s, 3H), 5.65 (d, 1H), 7.40 (m, 3H), 7.50 (m, 2H), 9.45 (s, 1H) ppm.

Example 79

(1-methyl-7-oxo-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-5-yl)(phenyl)-methanone

[0238]



[0239] Pyridinium chlorochromate (400mg, 0.00186mol) was added to a solution of 5-[α -hydroxybenzyl]-1-methyl-3-propyl-6,7-dihydro-1H-pyrazolo[4,3-d]pyrimidin-7-one (200mg, 0.00068mol) in dichloromethane (50ml) and the reaction stirred at room temperature for 3 hours.

[0240] The reaction mixture was filtered through silica gel, eluting with dichloromethane:methanol (95:5 by volume). Evaporation under reduced pressure of the desired fractions followed by crystallisation from ethyl acetate/hexane, gave the title compound as yellow needles (110mg), m.p.172-173°C.

[0241] Found: C, 64.81; H, 5.56; N, 18.95; $\text{C}_{16}\text{H}_{16}\text{N}_4\text{O}_2$, requires C, 64.85; H, 5.44; N, 18.91%.


[0242] $^1\text{H-NMR}$ (CDCl_3): δ = 1.05 (t, 3H), 1.90 (m, 2H), 2.95 (t, 2H), 4.35 (s, 3H), 7.55 (m, 2H), 7.70 (m, 1H), 8.50 (d, 2H) 10.15 (s, 1H) ppm.

SYNTHESIS PREPARATIONS

Preparation 1

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-bromophenyl)acetamide

[0243]



[0244] Oxalyl chloride (7.5ml, 0.86mol) was added dropwise to a solution of 4-bromophenylacetic acid (8.5g, 0.040mol) and dimethylformamide (3 drops) in dichloromethane and the reaction stirred at room temperature for 4 hours. The reaction mixture was then concentrated under reduced pressure and azeotroped with dichloromethane (30ml).

[0245] A solution of this 4-bromophenylacetyl chloride in dry dichloromethane (5ml) was then added dropwise to a solution of 4-amino-1-methyl-3-propyl-1H-5-pyrazolecarboxamide (6g, 0.033mol) and triethylamine (7ml, 0.050mol) in dichloromethane (250ml), and the reaction stirred at room temperature for 18 hours.


[0246] The reaction mixture was concentrated under reduced pressure and the residue triturated with 1N aqueous hydrochloric acid solution. The resulting precipitate was filtered and washed with water (50ml) and ether (50ml). Recrystallisation from ethanol gave the title compound as a solid (6.48g), m.p.241-243°C.

[0247] Found: C, 50.38; H, 5.00; N, 14.59%, $C_{16}H_{19}N_4BrO_2$, requires C, 50.67; H, 5.05; N, 14.77%.




[0248] 1H -NMR (DMSO- d_6): δ = 0.78 (t, 3H), 1.40 (m, 2H), 2.27 (t, 2H), 3.62 (s, 2H), 3.86 (s, 3H), 7.19 (s, 1H), 7.28 (d, 2H), 7.52 (d, 2H), 7.70 (s, 1H), 9.44 (s, 1H) ppm.




Preparations 2 to 22



[0249] The compounds of the following tabulated preparations of the general formula:







were prepared from 4-amino-1-methyl-3-propyl-1H-5-pyrazolecarboxamide and the appropriate acid chloride using similar methods to that described in Preparation 1.



Prep.No	R ₁	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting point/Crystallisation solvent
2	-CH ₃		<p>Found: C, 64.01; H, 6.72; N, 18.35%</p> <p>C₁₆H₂₀N₄O₂, requires C, 63.98; H, 6.71; N, 18.65%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.76 (t, 3H), 1.40 (m, 2H), 2.26 (t, 2H), 3.60 (s, 2H), 3.87 (s, 3H), 7.20-7.34 (m, 6H), 7.70 (s, 1H), 9.45 (s, 1H) ppm.</p> <p>Melting point: 238-240°C.</p> <p>Crystallisation solvent: ethanol.</p>
3	-CH ₂ CH ₃		<p>Found: C, 65.00; H, 7.04; N, 17.76%</p> <p>C₁₇H₂₂N₄O₂, requires C, 64.94; H, 7.05; N, 17.82%.</p> <p>¹H-NMR (CDCl₃): δ = 0.72 (t, 3H), 1.24 (t, 3H), 1.34 (m, 2H), 2.27 (t, 2H), 3.58 (s, 2H), 4.24 (q, 2H), 6.02 (s, 1H), 7.20 (m, 5H), 8.58 (s, 1H) ppm.</p> <p>Melting point: 191-192°C.</p> <p>Crystallisation solvent: ethanol.</p>
4	-CH ₃		<p>Found: C, 64.76; H, 7.05; N, 17.62%</p> <p>C₁₇H₂₂N₄O₂, requires C, 64.94; H, 7.05; N, 17.82%.</p> <p>¹H-NMR (CDCl₃): δ = 0.82 (t, 3H), 1.40 (m, 2H), 1.66 (d, 3H), 2.24 (t, 2H), 3.82 (q, 1H), 4.00 (s, 3H), 5.52 (s, 1H), 6.60 (s, 1H), 7.28 (s, 1H), 7.40 (m, 5H) ppm.</p> <p>Melting point: 213-215°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol</p>



5	-CH ₃		<p>Found: C, 65.79; H, 7.39; N, 16.85 %</p> <p>C₁₈H₂₄N₄O₂, requires C, 65.83; H, 7.37; N, 17.06 %.</p> <p>¹H-NMR (CDCl₃): d = 0.84 (t, 3H), 1.42 (m, 2H), 1.72 (s, 6H), 2.25 (t, 2H), 4.00 (s, 2H), 6.05 (s, 1H), 6.42 (s, 1H), 7.42 (m, 6H) ppm</p> <p>Melting point: 192-193°C.</p> <p>Crystallisation solvent: ethanol.</p>
6	-CH ₃		<p>Found: C, 65.87; H, 7.37; N, 17.05 %</p> <p>C₁₈H₂₄N₄O₂, requires C, 65.83; H, 7.37; N, 17.06 %.</p> <p>¹H-NMR (CDCl₃): d = 0.70 (t, 3H), 0.88 (t, 3H), 1.34 (m, 2H), 1.76 (m, 1H), 2.13 (m, 1H), 2.22 (t, 2H), 3.50 (t, 1H), 3.92 (s, 3H), 5.79 (s, 1H), 7.20-7.35 (m, 6H), 8.57 (s, 1H) ppm.</p> <p>Melting point: 226-227°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol</p>
7	-CH ₃		<p>Found: C, 66.42; H, 6.79; N, 17.39 %.</p> <p>C₁₈H₂₂N₄O₂, requires C, 66.24; H, 6.79; N, 17.17 %</p> <p>¹H-NMR (CDCl₃): d = 0.85 (t, 3H), 1.28 (m, 2H), 1.45 (m, 2H), 1.72 (m, 2H), 2.34 (t, 2H), 4.00 (s, 3H), 5.60 (s, 1H), 6.62 (s, 1H), 7.50 (m, 5H), 7.89 (s, 1H) ppm.</p> <p>Melting point: 176-177°C.</p> <p>Crystallisation solvent: ethanol.</p>



8	-CH ₃		<p>Found: C, 62.14; H, 6.71; N, 16.78%</p> <p>C₁₇H₂₂N₄O₃, requires C, 61.80; H, 6.71; N, 16.96%.</p> <p>¹H-NMR (CDCl₃): δ = 0.82 (t, 3H), 1.46 (m, 2H), 2.30 (t, 2H), 3.78 (s, 2H), 3.94 (s, 3H), 4.00 (s, 3H), 5.64 (s, 1H), 7.04 (m, 2H), 7.35 (m, 2H), 7.50 (s, 1H) ppm.</p> <p>Melting point: 208-209°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol</p>
9	-CH ₃		<p>Found: C, 61.43; H, 6.57; N, 16.94%</p> <p>C₁₇H₂₂N₄O₃, requires C, 61.80; H, 6.71; N, 16.96%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.77 (t, 3H), 1.40 (m, 2H), 2.26 (t, 2H), 3.54 (s, 2H), 3.73 (s, 3H), 3.86 (s, 3H), 6.88 (d, 2H), 7.22 (m, 3H), 7.70 (s, 1H), 9.39 (s, 1H) ppm.</p> <p>Melting point: 214-216°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol.</p>




10	-CH ₃		Found: C, 55.31, H, 5.59; N, 20.49% C ₁₆ H ₁₉ N ₅ O ₄ , requires C, 55.64; H, 5.55; N, 20.28%. ¹ H-NMR (DMSO-d ₆): d = 0.98 (t, 3H), 1.53 (m, 2H), 2.35 (t, 2H), 3.88 (s, 3H), 4.12 (s, 2H), 7.11 (s, 1H), 7.5 (m, 2H), 7.72 (m, 2H), 8.06 (d, 1H), 9.50 (s, 1H) ppm. Melting point: 229-231°C.
11	-CH ₃		Found: C, 55.82; H, 5.60; N, 20.53% C ₁₆ H ₁₉ N ₅ O ₄ , requires C, 55.64; H, 5.55; N, 20.28%. ¹ H-NMR (DMSO-d ₆): d = 0.75 (t, 3H), 1.41 (m, 2H), 2.28 (t, 2H), 3.82 (s, 2H), 3.86 (s, 3H), 7.20 (s, 1H), 7.66 (m, 2H), 7.78 (d, 1H), 8.14 (d, 1H), 8.12 (s, 1H), 9.57 (s, 1H) ppm. Melting point: 238-240°C.

12	-CH ₃		<p>Found: C, 55.97; H, 5.60; N, 20.10%</p> <p>C₁₆H₁₉N₅O₄, requires C, 55.64; H, 5.55; N, 20.28%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.78 (t, 3H), 1.42 (m, 2H), 2.28 (t, 2H), 3.82 (s, 2H), 3.86 (s, 3H), 7.18 (s, 1H), 7.60 (d, 2H), 7.70 (s, 1H), 8.20 (d, 2H), 9.57 (s, 1H) ppm.</p> <p>Melting point: 261-263°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol.</p>
13	-CH ₃		<p>Found: C, 50.59; H, 4.99; N, 14.85%</p> <p>C₁₆H₁₉N₄BrO₂, requires C, 50.67; H, 5.05; N, 14.77%</p> <p>¹H-NMR (DMSO-d₆): δ = 0.85 (t, 3H), 1.52 (m, 2H), 2.36 (t, 2H), 3.82 (s, 2H), 3.88 (s, 3H), 7.20 (m, 2H), 7.34 (m, 1H), 7.41 (d, 1H), 7.60 (d, 1H), 7.76 (s, 1H), 9.45 (s, 1H) ppm.</p> <p>Melting point: 250-252°C.</p> <p>Crystallisation solvent: acetonitrile.</p>

14	-CH ₃		<p>Found: C, 50.49; H, 5.04; N, 14.99%</p> <p>C₁₆H₁₉N₄BrO₂, requires C, 50.67; H, 5.05; N, 14.77%</p> <p>¹H-NMR (DMSO-d₆): δ = 0.87 (t, 3H), 1.41 (m, 2H), 2.26 (t, 2H), 3.62 (s, 2H), 3.86 (s, 3H), 7.21 (s, 1H), 7.30 (m, 2H), 7.46 (m, 1H), 7.72 (s, 1H), 9.47 (s, 1H) ppm.</p> <p>Melting point: 240-242°C.</p> <p>Crystallisation solvent: acetonitrile.</p>
15	-CH ₃		<p>Found: C, 51.78; H, 5.40; N, 14.15%</p> <p>C₁₇H₂₁N₄BrO₂, requires C, 51.92; H, 5.38; N, 14.25%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.70 (t, 3H), 1.30 (m, 2H), 1.38 (d, 3H), 2.16 (t, 2H), 3.86 (s, 3H), 7.12 (s, 1H), 7.73 (d, 2H), 7.52 (d, 2H), 7.69 (s, 1H), 9.35 (s, 1H) ppm.</p> <p>Melting point: 234-236°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol.</p>

16	-CH ₃		<p>¹H-NMR (DMSO-d₆): d = 0.76 (t, 3H), 1.40 (m, 2H), 2.28 (t, 2H), 3.62 (s, 2H), 3.88 (s, 3H), 7.18 (s, 1H), 7.35 (m, 4H), 7.70 (s, 1H), 9.46 (s, 1H) ppm.</p> <p>Crystallisation solvent: acetonitrile.</p>
17	-CH ₃		<p>Found: C, 55.65; H, 5.28; N, 14.94 %</p> <p>C₁₇H₁₉N₄F₃O₂, requires C, 55.43; H, 5.20; N, 15.21 %.</p> <p>¹H-NMR (DMSO-d₆): d = 0.87 (t, 3H), 1.32 (m, 2H), 2.34 (t, 2H), 3.86 (s, 3H), 3.92 (s, 2H), 7.18 (s, 1H), 7.50 (m, 2H), 7.70 (m, 3H), 9.45 (s, 1H) ppm.</p> <p>Melting point: 247-249°C.</p> <p>Crystallisation solvent: acetonitrile.</p>


18	-CH ₃		<p>Found: C, 55.63; H, 5.24; N, 15.11%</p> <p>C₁₇H₁₉N₄F₃O₂, requires C, 55.43; H, 5.20; N, 15.21%.</p> <p>¹H-NMR (DMSO-d₆): d = 0.76 (t, 3H), 1.39 (m, 2H), 2.26 (t, 2H), 3.75 (s, 2H), 3.86 (s, 3H), 7.22 (s, 1H), 7.55 (d, 2H), 7.70 (m, 3H), 9.52 (s, 1H) ppm.</p> <p>Melting point: 225-228°C.</p> <p>Crystallisation solvent: ethyl acetate/methanol.</p>
19	-CH ₃		<p>Found: C, 66.99; H, 7.47; N, 16.35%</p> <p>C₁₉H₂₆N₄O₂, requires C, 66.64; H, 7.65; N, 16.36%.</p> <p>¹H-NMR (DMSO-d₆): d = 0.75 (t, 3H), 1.18 (d, 6H), 1.38 (m, 2H), 2.25 (t, 2H), 2.86 (m, 1H), 3.57 (s, 2H), 3.86 (s, 3H), 7.22 (m, 5H), 7.72 (s, 1H), 9.42 (s, 1H) ppm.</p> <p>Melting point: 200-201°C.</p> <p>Crystallisation solvent: ethyl acetate</p>

20	-CH ₃		¹ H-NMR (DMSO-d ₆): d = 0.74 (t, 3H), 1.40 (m, 2H), 2.25 (t, 2H), 2.57 (s, 3H), 3.72 (s, 2H), 3.85 (s, 3H), 7.47 (d, 2H), 7.92 (d, 2H), 9.52 (s, 2H) ppm.
21	-CH ₃		Found: C, 45.35; H, 4.32; N, 16.26% C ₁₆ H ₁₈ N ₃ BrO ₄ , requires C, 45.29; H, 4.28; N, 16.57%. ¹ H-NMR (DMSO-d ₆): d = 0.88 (t, 3H), 1.52 (m, 2H), 2.33 (t, 2H), 3.86 (s, 3H), 4.08 (s, 2H), 7.08 (s, 1H), 7.54 (d, 1H), 7.73 (s, 1H), 7.92 (d, 1H), 8.24 (s, 1H), 9.49 (s, 1H) ppm. Melting point: 249-251°C. Crystallisation solvent: acetonitrile.
22	-CH ₃		Found: C, 52.02; H, 5.45; N, 14.48% C ₁₇ H ₂₁ N ₄ BrO ₂ , requires, C, 51.91; H, 5.38; N, 14.25%. ¹ H-NMR (DMSO-d ₆): d = 0.78 (t, 3H), 1.40 (m, 2H), 2.26 (t, 2H), 3.62 (s, 2H), 3.85 (s, 3H), 4.70 (s, 2H), 7.22 (s, 1H), 7.30 (d, 2H), 7.40 (d, 2H), 7.62 (s, 1H), 9.45 (s, 1H) ppm. Melting point: 191-193°C. Crystallisation solvent: acetonitrile.

Preparation 23

4-*tert*-butyl carbamoylmethoxyphenyl acetic acid methyl ester

[0250]



[0251] Methyl 4-hydroxyphenylacetate (4.1g, 0.0247mol) was added to a suspension of sodium hydride (800mg, 80%, 0.0266mol) in dimethylformamide (100ml) and the mixture stirred at room temperature for 30 minutes. *Tert*-butyl bromoacetate (4.2ml, 0.0258mol) was added dropwise and the resulting solution stirred for a further 2 hours. Water (250ml) and 1N aqueous hydrochloric acid (100ml) were then added and the mixture extracted with diethyl ether (2x250ml). The combined organic extracts were dried (MgSO_4), filtered and evaporated under reduced pressure.

[0252] Purification by flash column chromatography, eluting with dichloromethane gave the title compound as a colourless liquid (5.67g).


[0253] Found: C, 64.20; H, 7.13; $\text{C}_{15}\text{H}_{20}\text{O}_5$, requires C, 64.27; H, 7.13%

[0254] $^1\text{H-NMR}$ (CDCl_3): δ = 1.52 (s, 9H), 3.59 (s, 2H), 3.72 (s, 3H), 4.54 (s, 2H), 6.88 (d, 2H), 7.24 (d, 2H) ppm.

Preparation 24

4-(1-methyl piperidinoxy)-phenylacetic acid methyl ester

[0255]



[0256] Diethylazodicarboxylate (2.1ml, 0.013mol) was added dropwise to a solution of methyl 4-hydroxyphenylacetate (2.2g, 0.013mol), 4-hydroxy-1-methylpiperidine (1.5g, 0.013mol) and triphenylphosphine (3.5g, 0.0133mol) in tetrahydrofuran (50ml) and the reaction stirred at room temperature for 20 hours. The reaction mixture was then concentrated under reduced pressure.


[0257] Purification by flash column chromatography eluting with 0.880 aqueous ammonia:methanol:dichloromethane (0.5:5:95 by volume) gave the title compound as an oil (1.48g).

[0258] $^1\text{H-NMR}$ (CDCl_3): δ = 1.88 (m, 2H), 2.02 (m, 2H), 2.32 (m, 5H), 2.72 (m, 2H), 3.58 (s, 2H), 3.72 (s, 3H), 4.32 (m, 1H), 6.88 (d, 2H), 7.20 (d, 2H) ppm.

Preparation 25

4-*tert*-butyl carbamoylmethoxyphenyl acetic acid

[0259]



[0260] 1N aqueous sodium hydroxide solution (16ml) was added to a solution of 4-*tert*-butyl carbamoylmethoxyphenylacetic acid methyl ester (2.8g, 0.010mol) in methanol (10ml) and the reaction stirred at room temperature for 5 hours. The reaction mixture was then concentrated under reduced pressure, the residue suspended in 1N aqueous hydrochloric acid (20ml) and extracted with diethyl ether (2x25ml). The combined organic extracts were dried (MgSO₄), filtered and evaporated under reduced pressure.

[0261] Purification by flash column chromatography, eluting with a solvent gradient of diethyl ether:hexane (50:50 to 67:33 by volume) gave the title compound as a solid (680mg), m.p. 97-98°C.

[0262] Found: C, 62.74; H, 6.89; C₁₄H₁₈O₅, requires C, 63.14; H, 6.81%.

[0263] ¹H-NMR (CDCl₃): δ = 1.54 (s, 9H), 3.62 (s, 2H), 4.54 (s, 2H), 6.88 (d, 2H), 7.24 (d, 2H) ppm.

Preparation 26

4-(1-methyl piperidinoxy)-phenylacetic acid

[0264] The title compound was prepared using a similar method to that described in Preparation 25 from 4-(1-methyl piperidinoxy)-phenylacetic acid methyl ester and was obtained, after trituration with acetonitrile, as a solid (82%), m.p. 159-161°C.


[0265] Found: C, 67.48; H, 7.65; N, 5.57; C₁₄H₁₉NO₃, requires C, 67.44; H, 7.68; N, 5.62%.

[0266] ¹H-NMR (DMSO-d₆): δ = 1.62 (m, 2H), 1.88 (m, 2H), 2.18 (m, 5H), 2.60 (m, 2H), 3.48 (s, 2H), 4.30 (m, 1H), 6.84 (d, 2H), 7.15 (d, 2H) ppm.

Preparation 27

tert-butyl-2-(4-[[5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl]carbamoyl]methoxy)phenylacetate

[0267]



[0268] Phosphorous trichloride (141mg, 0.0010mol) was added dropwise to an ice-cooled solution of 4-amino-1-methyl-3-propyl-1H-5-pyrazolecarboxamide (370mg, 0.0020mol) in pyridine (6ml), and the reaction stirred for an hour at room temperature. 4-*tert*-butyl carbamoylmethoxy-phenyl acetic acid (600mg, 0.0023mol) was then added and stirring continued at reflux for 3 hours. On cooling, the mixture was concentrated under reduced pressure.

[0269] Purification by flash column chromatography eluting with a solvent gradient of methanol:dichloromethane (2:98 to 5:95 by volume), followed by crystallisation from ethyl acetate/hexane gave the title compound as a solid (465mg), m.p.145-146°C.

[0270] Found: C, 61.07; H, 6.97; N, 12.76; $C_{22}H_{30}N_4O_5$ requires C, 61.38; H, 7.02; N, 13.02%.

[0271] 1H -NMR ($CDCl_3$): δ = 0.88 (t, 3H), 1.52 (m, 11H), 2.34 (t, 2H), 3.74 (s, 2H), 4.01 (s, 3H), 4.57 (s, 2H), 5.58 (s, 2H), 6.70 (s, 1H), 6.97 (d, 2H), 7.29 (d, 2H) ppm.

Preparation 28

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-[4-(1-methylpiperidinoxy)acetamide

[0272] The title compound was prepared from 4-(1-methyl piperidinoxy)-phenylacetic acid following the procedure described in Preparation 27 and was obtained as a solid (47%), m.p. 170-171°C.


[0273] Found: C, 63.59; H, 7.62; N, 16.87; $C_{22}H_{31}N_5O_3$, requires C, 63.90; H, 7.56; N, 16.94%.

[0274] 1H -NMR ($CDCl_3$): δ = 0.87 (t, 3H), 1.48 (m, 2H), 1.88 (m, 2H), 2.04 (m, 2H), 2.33 (m, 7H), 2.72 (m, 2H), 3.73 (s, 2H), 4.00 (s, 3H), 4.36 (m, 1H), 5.56 (s, 1H), 6.68 (s, 1H), 6.98 (d, 2H), 7.26 (d, 2H), 7.54 (s, 1H) ppm.

Preparation 29

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(1,3-benzodioxol-5-yl)acetamide

[0275]



[0276] Oxalyl chloride (0.6ml, 0.0067mol) was added dropwise to a solution of 3,4-methylenedioxyphenylacetic acid (600mg, 0.0033mol) and dimethylformamide (1 drop) in dichloromethane and the reaction stirred at room temperature for 2 hours. The reaction mixture was concentrated under reduced pressure, and azeotroped with dichloromethane (40ml).

[0277] This acid chloride was added to a solution of 4-amino-1-methyl-3-propyl-1H-5-pyrazolecarboxamide (550mg, 0.0030mol) in pyridine (6ml) and the reaction stirred at room temperature for 18 hours. The reaction mixture was then concentrated under reduced pressure.

[0278] Purification by flash column chromatography, eluting with a solvent gradient of methanol:dichloromethane (2:98 to 5:95 by volume), followed by crystallisation from acetonitrile gave the title compound (420mg), m.p.231-233°C.

[0279] Found: C, 59.25; H, 5.79; N, 16.62. $C_{17}H_{20}N_4O_4$, requires C, 59.29; H, 5.85; N, 16.27%.

[0280] 1H -NMR ($DMSO-d_6$): δ = 0.78 (t, 3H), 1.40 (m, 2H), 2.26 (t, 2H), 3.52 (s, 2H), 3.86 (s, 3H), 5.98 (s, 2H), 6.78 (s, 1H), 6.86 (d, 2H), 7.18 (s, 1H), 7.70 (s, 1H), 9.37 (s, 1H) ppm.

Preparation 30

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-phenoxyphenyl-5-yl)acetamide

[0281] The title compound was prepared from 4-phenoxyphenylacetic acid and 4-amino-1-methyl-3-propyl-1H-5-pyrazolecarboxamide following the procedure described in Preparation 29 and was obtained as a solid (21%), m.p. 192-194°C.


[0282] Found: C, 67.28; H, 6.20; N, 14.49; $C_{22}H_{24}N_4O_3$, requires C, 67.33; H, 6.16; N, 14.28%.

[0283] 1H -NMR ($CDCl_3$): δ = 0.89 (t, 3H), 1.52 (m, 2H), 2.36 (t, 2H), 3.78 (s, 2H), 4.02 (s, 3H), 5.57 (s, 1H), 6.71 (s, 1H), 7.08 (m, 4H), 7.18 (m, 1H), 7.36 (m, 5H) ppm.

Preparation 31

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-[4-(morpholinomethyl)phenyl]acetamide

[0284]



[0285] Morpholine (0.3ml, 0.00344mol) was added to a solution of N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-bromobenzyl)acetamide (400mg, 0.001mol) in acetonitrile (3ml) and the reaction stirred at reflux for 2 hours. On cooling, the reaction was concentrated under reduced pressure.


[0286] Purification by flash column chromatography, eluting with a solvent gradient of 0.880 aqueous ammonia: methanol:dichloromethane (0.5:95 to 0.5:5:95 by volume), followed by crystallisation from ethyl acetate gave the title compound (350mg), m.p. 148-150°C.

[0287] Found: C, 63.48; H, 7.36; N, 17.16; $C_{21}H_{29}N_5O_3$, requires C, 63.12; H, 7.32; N, 17.53%.



[0288] $^1\text{H-NMR}$ (CDCl_3): δ = 0.85 (t, 3H), 1.48 (m, 2H), 2.32 (t, 2H), 2.47 (m, 4H), 3.54 (s, 2H), 3.74 (m, 4H), 3.80 (s, 2H), 4.00 (s, 3H), 5.58 (s, 1H), 6.68 (s, 1H), 7.32 (d, 2H), 7.44 (d, 2H), 7.50 (s, 1H) ppm.

Preparation 32 and 33

[0289] The compounds of the following tabulated preparations of the general formula:




were prepared from N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-bromobenzyl)acetamide and the appropriate amine, using similar methods to that described in Preparation 31.

Prep.No	Ar(R ₂)(R ₃)C-	Analysis/ ¹ H-NMR/Melting point/Crystallisation solvent
32		<p>Found: C, 63.81; H, 7.64; N, 19.74% C₁₉H₂₇N₅O₂, requires C, 63.84; H, 7.61; N, 19.59%.</p> <p>¹H-NMR (CDCl₃): δ = 0.86 (t, 3H), 1.48 (m, 2H), 2.28 (m, 8H), 2.32 (t, 2H), 3.47 (s, 2H), 4.00 (s, 3H), 5.58 (s, 1H), 6.71 (s, 1H), 7.32 (d, 2H), 7.40 (d, 2H), 7.50 (s, 1H) ppm.</p> <p>Melting point: 184-186°C. Crystallisation solvent: ethyl acetate/hexane.</p>
33		<p>Found: C, 63.23; H, 6.24; N, 22.27% C₂₀H₂₄N₆O₂, requires C, 63.14; H, 6.36; N, 22.09%.</p> <p>¹H-NMR (DMSO-d₆): δ = 0.74 (t, 3H), 1.38 (m, 2H), 2.24 (t, 2H), 3.60 (s, 2H), 3.84 (s, 3H), 5.16 (s, 2H), 6.88 (s, 1H), 7.16 (s, 1H), 7.20 (d, 2H), 7.30 (d, 2H), 7.72 (s, 1H), 9.42 (s, 1H) ppm.</p> <p>Melting point: 222-224°C. Crystallisation solvent: ethanol.</p>

Preparation 34

N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-[4-(ethoxymethyl)phenyl]acetamide

[0290]



[0291] Silver nitrate (290mg, 0.0017mol) was added to a solution of N-(5-carbamoyl-1-methyl-3-propyl-1H-4-pyrazolyl)-2-(4-bromophenyl)acetamide (600mg, 0.0015mol) in ethanol (5ml) and the reaction stirred at reflux for 5 hours

and then at room temperature for 18 hours.


[0292] Ethanol (10ml) was added, the mixture filtered, and the solid washed with further ethanol.

[0293] The filtrate was evaporated under reduced pressure and the residue crystallised from ethanol to give the title compound, as a solid (500mg).

Preparation 35

2-(4-bromophenyl)propionic acid

[0294]




[0295] The title compound was prepared following a similar procedure to that described in *Synthesis* 1982; 456.

Preparation 36

4-acetyl phenylacetic acid

[0296]




[0297] The title compound was prepared following a similar procedure to that described in *J.A.C.S.* 1946; 68; 2133.

Preparation 37

2-nitro-4-bromophenylacetic acid

[0298]




[0299] The title compound was prepared following a similar procedure to that described in *Chem. Pharm. Bull.*; 1985; 33; 1414.

Preparation 38

3-pyridinesulphonyl chloride hydrochloride


[0300]

[0301] The title compound was prepared following a similar procedure to that described in *Annalen*; 1939; 72; 77.

Preparation 39

2-(methylaminomethyl)pyridine

[0302]



[0303] The title compound was prepared following a similar procedure to that described in US-A-2798075.

ACTIVITY STUDIES

[0304] Initially there is presented a Protocol for measuring PDE inhibitory activity.

PROTOCOLPhosphodiesterase (PDE) inhibitory activity

[0305] *In vitro* PDE inhibitory activities against cyclic guanosine 3',5'-monophosphate (cGMP) and cyclic adenosine 3',5'-monophosphate (cAMP) phosphodiesterases can be determined by measurement of their IC₅₀ inhibition of enzyme activity.

[0306] The required PDE enzymes are isolated from a variety of sources, including rat kidney, human corpus cavernosum, human platelets, rabbit platelets, human cardiac ventricle, human skeletal muscle and bovine retina, essentially by the method of W J Thompson and M M Appleman (Biochem, 1971, 10, 311).

[0307] For example, for some of the studies the calcium/calmodulin (Ca/CAM)-dependent PDE (PDE1) is obtained from either human cardiac ventricle or rat kidney. The cGMP-stimulated PDE (PDE2), the cGMP-specific PDE (PDE5) and the cGMP-inhibited cAMP PDE (PDE3) are obtained from human corpus cavernosum tissue. PDE5 is also obtained from human platelets or rabbit platelets by techniques usual in the art. The cAMP-specific PDE (PDE4) is obtained from rat kidney. The photoreceptor PDE (PDE6) is obtained from bovine retina.


[0308] Assays are performed using a modification of the "batch" method of W J Thompson *et al* (Biochem, 1979, 18, 5228).

RESULTS

[0309] The compounds of the present invention were tested for PDE inhibition. The results showed that the compounds are inhibitors of at least Ca/CAM-dependent PDE1. Some of the compounds are selective and potent inhibitors of Ca/CAM-dependent PDE1.

[0310] In particular, we found the following results for a preferred compound of the present invention having the

formula:



which for ease of reference is referred to as the Example 50 compound.

Example 50		
PDE TYPE	SOURCE	IC ₅₀
1	Human cardiac ventricle	38nM
2	Human corpus cavernosum	1.99μM
3	Human corpus cavernosum	3.94μM
4	Rat Kidney	23μM
5	Human corpus cavernosum	2.49μM
6	Bovine retina	2.03μM
(IC denotes inhibitory concentration)		

COMPARATIVE ACTIVITY STUDIES

[0311] For these studies we compared the activity of the '188 compound with the Example 50 compound.


[0312] The results are as follows:

	cGMP / PDE1 INHIBITION - IC ₅₀ Values	
SOURCE	EXAMPLE 50	'188 COMPOUND
Rat Kidney	37nM	9.9μM


	PDE5 INHIBITION - IC ₅₀ Values	
SOURCE	EXAMPLE 50	'188 COMPOUND
Rabbit Platelet	6.7μM	2.8μM
Human Platelet	2.7μM	3.2μM

FURTHER STUDIES

[0313] In addition, we investigated the following two compounds.



15 (which for ease of reference is referred to as the Example 36 compound)



30 (which for ease of reference is referred to as the Example 37 compound)

[0314] The results of these additional studies are presented below.

35

EXAMPLE No	RAT KIDNEY PDE1	RABBIT PLATELET PDE5
EXAMPLE 36	59nM	1.8μM
EXAMPLE 37	94nM	6.4μM


SUMMARY OF RESULTS

40 [0315] The results demonstrate that the compounds of the present invention - especially the Example 50, the Example 36 and the Example 37 compounds presented above - are potent and selective PDE1 inhibitors.

[0316] Other modifications will be apparent to those skilled in the art.

Claims

- 45
1. A compound of the formula (I)
- 50
- 55



15 wherein

R_6 is C_2 - C_6 alkyl;

R_1 is H or C_1 - C_4 alkyl;


20 each of R_2 and R_3 is independently selected from H and C_1 - C_4 alkyl, or

R_2 is H or C_1 - C_4 alkyl and R_3 is OH, C_2 - C_4 alkanoyloxy or fluoro, or

R_2 and R_3 when taken together represent C_2 - C_6 alkylene, or

R_2 and R_3 when taken together with the carbon atom to which they are attached represent a carbonyl group;

25 Ar is either (a)



40 wherein

each of R_4 , R_5 and R_6 is independently selected from

45 H,

C_1 - C_4 alkyl,

C_1 - C_4 alkoxy,

C_1 - C_4 alkoxy-Z-,

halo,

halo(C_1 - C_4)alkyl,

50 phenoxy, optionally substituted by up to three substituents each of which substituent is independently selected from halo, C_1 - C_4 alkyl, and C_1 - C_4 alkoxy,

nitro,

hydroxy,

hydroxy-Z-,

55 C_2 - C_4 alkanoyl,

amino,

amino-Z-,

(C_1 - C_4 alkyl)NH,

(C₁-C₄ alkyl)₂N-,
 (C₁-C₄ alkyl)NH-Z-,
 (C₁-C₄ alkyl)₂N-Z-,
 -COOH,
 -Z-COOH,
 -COO(C₁-C₄ alkyl),
 -Z-COO(C₁-C₄ alkyl)
 C₁-C₄ alkanesulphonamido,
 C₁-C₄ alkanesulphonamido-Z-,
 halo(C₁-C₄)alkanesulphonamido,
 halo(C₁-C₄)alkanesulphonamido-Z-,
 C₁-C₄ alkanamido,
 C₁-C₄ alkanamido-Z-,
 HOOC-Z-NH-,
 HOOC-Z-NH-Z-,
 (C₁-C₄ alkyl)OOC-Z-NH-,
 (C₁-C₄ alkyl)OOC-Z-NH-Z-,
 C₁-C₄ alkyl-NH-SO₂-NH-,
 C₁-C₄ alkyl-NH-SO₂-NH-Z-,
 (C₁-C₄ alkyl)₂-N-SO₂-NH-,
 (C₁-C₄ alkyl)₂-N-SO₂-NH-Z-,
 C₁-C₄ alkoxy CH=CH-Z-CONH-,
 C₁-C₄ alkoxy CH=CHCONH
 C₁-C₄ alkyl-SO₂-N(C₁-C₄ alkyl)-,
 C₁-C₄ alkyl-SO₂-N(C₁-C₄ alkyl)-Z-,
 (C₁-C₄ alkyl)NH-Z-SO₂-NH-,
 (C₁-C₄ alkyl)₂N-Z-SO₂-NH-,
 (C₁-C₄ alkyl)NH-Z-SO₂-NH-Z-,
 (C₁-C₄ alkyl)₂N-Z-SO₂-NH-Z-,
 benzenesulphonamido, optionally ring substituted by up to three substituents each of which is independently selected from halo, C₁-C₄ alkyl, and C₁-C₄ alkoxy,
 C₁-C₄ alkanoyl-N(C₁-C₄ alkyl)-,
 C₁-C₄ alkanoyl-N(C₁-C₄ alkyl)-Z-,
 C₁-C₄ alkoxy carbonyl-CH(CH₂OH)NHSO₂-,
 -SO₃H,
 -SO₂NH₂,
 H₂NOC-CH(CH₂OH)-NHSO₂-,
 HOOC-Z-O-, and
 (C₁-C₄ alkyl)OOC-Z-O-,

or optionally one of R₄, R₅ and R₆ is a G-Het group and wherein the others of R₄, R₅ and R₆ are independently selected from the R₄, R₅ and R₆ substituents listed above;

Z is C₁-C₄ alkylene,

G is a direct link, Z, O, -SO₂NH-, SO₂, or -Z-N(C₁-C₄ alkyl)SO₂-,

Het is a 5- or 6-membered heterocyclic group containing 1, 2, 3 or 4 nitrogen heteroatoms; or 1 or 2 nitrogen heteroatoms and 1 sulphur heteroatom or 1 oxygen heteroatom; or the heterocyclic group is furanyl or thiophenyl; wherein the Het group is saturated or partially or fully unsaturated and optionally substituted by up to 3 substituents, wherein each substituent is independently selected from C₁-C₄ alkyl, oxo, hydroxy, halo, and halo(C₁-C₄) alkyl;

or (b) any one of the following bicyclic groups:

benzodioxolanyl,
 benzodioxanyl,
 benzimidazolyl,

quinoliny,
indolyl,
quinazoliny,
isoquinoliny,
benzotriazolyl,
benzofuranyl,
benzothiophenyl,
quinoxaliny, or
phthaliziny,

wherein said bicyclic Ar groups are linked to the neighbouring -C(R₂R₃)- group *via* the benzo ring portion,

and wherein the heterocyclic portion of said bicyclic Ar group is optionally partially or fully saturated, said group being optionally substituted by one or more of C₁-C₄ alkyl, halo, hydroxy, oxo, amino, and C₁-C₄ alkoxy;

or a pharmaceutically acceptable salt of the compound, or a pharmaceutically acceptable solvate of the compound or the salt.

2. A compound, salt or solvate according to claim 1 wherein R_a is a C₂₋₅ alkyl group.

3. A compound, salt or solvate according to claim 1 or claim 2 wherein R_a is a C₂₋₄ alkyl group.

4. A compound, salt or solvate according to any one of claims 1 to 3 wherein R_a is a C₃ alkyl group.

5. A compound, salt or solvate according to any one of the preceding claims wherein R₁ is a C₁₋₆ alkyl group.

6. A compound, salt or solvate according to any one of the preceding claims wherein R₁ is a C₁₋₃ alkyl group.

7. A compound, salt or solvate according to any one of the preceding claims wherein R₁ is a methyl group.

8. A compound, salt or solvate according to any one of the preceding claims wherein R₂ is H.

9. A compound, salt or solvate according to any one of the preceding claims wherein R₃ is H.

10. A compound, salt or solvate according to any one of the preceding claims wherein R₄, R₅ and R₆ are independently selected from H, (C₁₋₄ alkyl)₂N-, C₁₋₄ alkanesulphonamido and benzenesulphonamido.

11. A compound, salt or solvate according to any one of the preceding claims wherein R₄, R₅ and R₆ are independently selected from H, diethylamino, methanesulphonamido and benzenesulphonamido.

12. A compound, salt or solvate according to any one of the preceding claims wherein Ar is 4-diethylaminophenyl.


13. A compound, salt or solvate according to any one of claims 1 to 11 wherein Ar is 2-methanesulphonamidophenyl.

14. A compound, salt or solvate according to any one of claims 1 to 11 wherein Ar is 4-benzenesulphonamidophenyl.


15. A compound, salt or solvate according to any one of claims 1 to 10 wherein one of R₄, R₅ and R₆ is (C₁₋₄ alkyl)₂N- and wherein the other two of R₄, R₅ and R₆ are H.

16. A compound, salt or solvate according to claim 15 wherein one of R₄, R₅ and R₆ is diethylamino and wherein the other two of R₄, R₅ and R₆ are H.


17. A compound, salt or solvate according to claim 1 wherein the compound is of the formula:



15 18. A compound, salt or solvate according to claim 1 wherein the compound is of the formula:



30 19. A compound, salt or solvate according to claim 1 wherein the compound is of the formula:



45 20. A pharmaceutical composition comprising a compound, salt or solvate according to any one of the preceding claims admixed with a pharmaceutically acceptable carrier, diluent or excipient.

21. A veterinary composition comprising a compound, salt or solvate according to any one of claims 1 to 19 admixed with a veterinarily acceptable carrier, diluent or excipient.

50 22. A compound, salt or solvate according to any one of claims 1 to 19 for use in medicine.

23. Use of a compound, salt or solvate according to any one of claims 1 to 19 in the manufacture of a pharmaceutical composition to inhibit PDE1 activity.

55 24. Use of a compound, salt or solvate according to any one of claims 1 to 19 in the manufacture of a veterinary composition to inhibit PDE1 activity.

25. A method of treatment to inhibit PDE1 activity comprising administering to a subject in need of treatment a compound, salt or solvate according to any one of claims 1 to 19 or a composition according to claim 20 or claim 21, wherein PDE1 activity is inhibited.

26. Use of a compound, salt or solvate according to any one of claims 1 to 19 in the manufacture of a pharmaceutical composition to treat stroke, dementia, memory enhancement, atherosclerosis, urge incontinence, hypertension, angina pectoris, congestive heart failure, myocardial infarction or restenosis.

27. Use of a compound, salt or solvate according to any one of claims 1 to 19 for the manufacture of a human medicament for the treatment of a medical condition for which a PDE1 inhibitor is indicated.

28. Use of a compound, salt or solvate according to any one of claims 1 to 19 for the manufacture of an animal medicament for the treatment of a medical condition for which a PDE1 inhibitor is indicated.

29. Use of a compound, salt or solvate according to any one of claims 1 to 19 for the manufacture of a human medicament for the treatment of male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.

30. Use of a compound, salt or solvate according to any one of claims 1 to 19 for the manufacture of an animal medicament for the treatment of male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility.

31. A method of treating or preventing a medical condition for which a PDE1 inhibitor is indicated, in a mammal (including a human being), which comprises administering to said mammal a therapeutically effective amount of a compound, salt or solvate according to any one of claims 1 to 19, or a pharmaceutical composition or veterinary formulation containing any of the foregoing.

32. A method of treating male erectile dysfunction, female sexual dysfunction, premature labour, dysmenorrhoea, benign prostatic hyperplasia (BPH), bladder outlet obstruction, incontinence, stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, stroke, peripheral vascular disease, conditions of reduced blood vessel patency, chronic asthma, bronchitis, allergic asthma, allergic rhinitis, glaucoma or diseases characterised by disorders of gut motility in a mammal (including a human being), which comprises administering to said mammal a therapeutically effective amount of a compound, salt or solvate according to any one of claims 1 to 19, or a pharmaceutical composition or veterinary formulation containing any of the foregoing.

33. A method of treating stroke, dementia, memory enhancement, atherosclerosis, urge incontinence, hypertension, angina pectoris, congestive heart failure, myocardial infarction or restenosis, which comprises administering to a subject a therapeutically effective amount of a compound, salt or solvate according to any one of claims 1 to 19, or a pharmaceutical composition or veterinary formulation containing any of the foregoing.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention shall be considered, for the purposes of subsequent proceedings, as the European search report

EP 98 30 8177

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
D,A	EP 0 201 188 A (WARNER-LAMBERT) 17 December 1986 * column 9, line 45 - column 14, line 45; claim 1 * -----	1, 20, 23	C07D487/04 A61K31/505 //(C07D487/04, 239:00,231:00)
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			C07D A61K
INCOMPLETE SEARCH			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>Although claims 25 and 31 to 32 are directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.</p>			
Place of search		Date of completion of the search	Examiner
THE HAGUE		27 January 1999	Alfaro Faus, I
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO FORM 1503 03 02 (P04C07)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 98 30 8177

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

27-01-1999

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 201188 A	17-12-1986	US 4666908 A JP 61236778 A	19-05-1987 22-10-1986

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82